

Kobayashi length bounds on bordered surfaces and generalized integral points on abelian varieties

Paolo Dolce

Abstract

Let B be a compact Riemann surface and $B_0 \subset B$ a bordered hyperbolic subsurface obtained by removing finitely many disjoint closed disks. Fix a nontrivial loop α in B_0 . For $s \geq 0$, let $L(\alpha, s)$ denote the supremum, over all finite subsets $S \subset B_0$ with $\#S \leq s$, of the minimal Kobayashi length of a loop in $B_0 \setminus S$ that is freely homotopic to α in B_0 . Phung in [10] proved that $L(\alpha, s)$ grows at most linearly and at least as $\sqrt{s}/\log s$. We sharpen the upper bound to $O(\sqrt{s \log s})$, which determines $\lim_{s \rightarrow \infty} \frac{\log L(\alpha, s)}{\log s} = \frac{1}{2}$, answering a question raised in [10, Question 1.4]. As an application, we improve the counting bound for generalized integral points on abelian varieties over complex function fields: for an abelian variety of dimension n over $\mathbb{C}(B)$, Phung proved that the number of (s, B_0) -generalized integral points modulo the constant trace grows at most as s^{2nk} , where $k = \text{rk}(\pi_1(B_0))$. We sharpen this to $s^{nk+\varepsilon}$ for every $\varepsilon > 0$, halving the exponent.

1	Introduction	1
1.1	Generalized integral points and the Lang–Vojta conjecture	1
1.2	Main results	2
2	Kobayashi length estimates	5
2.1	Kobayashi pseudo-metric	5
2.2	Growth of loop lengths in terms of the cusps	6
2.3	Reduction to the simple case	12
3	Bounding generalized integral points	13
3.1	Phung’s bound	13
3.2	Improved bound	18
	References	19

1 Introduction

1.1 Generalized integral points and the Lang–Vojta conjecture

Let k be an algebraically closed field of characteristic 0 and let B be a non-singular, irreducible, projective algebraic curve over k with function field $K = k(B)$. Let X be a smooth, integral, projective K -variety and fix a reduced effective divisor $D \subset X$. A *model* of (X, D) is a pair $(\mathcal{X}, \mathcal{D})$ where $\mathcal{X}$ is a normal k -variety equipped with a proper, flat morphism $f: \mathcal{X} \rightarrow B$ satisfying $\mathcal{X}_K \cong X$, and $\mathcal{D} \subset \mathcal{X}$ is a horizontal divisor with $\mathcal{D}_K \cong D$. Such a model always exists. We also define

$$\text{Sing}(f) := \{b \in B : \mathcal{X}_b \text{ is singular}\}.$$

The set of rational points $X(K)$ corresponds bijectively to the sections of f : for any $P \in X(K)$, let $\sigma_P: B \rightarrow \mathcal{X}$ be the section obtained as the Zariski closure of P in $\mathcal{X}$. By abuse of notation we will simply write $\sigma_P \in X(K)$. Given a finite set $S \subset B$, the set of $(S, \mathcal{D})$ -integral points of $\mathcal{X}$ is

$$\mathcal{X}(\mathcal{O}_{S, \mathcal{D}}) := \{\sigma_P \in X(K) : f(\sigma_P(B) \cap \mathcal{D}) \subseteq S\},$$

that is, the set of sections whose image doesn’t intersect the divisor $\mathcal{D}$ “above” $B \setminus S$.

Recall that the pair (X, D) is said to be of *log general type* when D is a normal crossings divisor and the log canonical bundle $\omega_X(D)$ is big. A special case is when $X = A$ is an abelian variety: the canonical bundle ω_A is trivial, so the log canonical bundle $\omega_A(D) \cong \mathcal{O}_A(D)$ is big whenever D is big.

The *geometric Lang-Vojta conjecture* predicts strong constraints for the integral points of pairs of log general type

Conjecture 1.1 (Geometric Lang-Vojta). Let (X, D) be a pair of log general type and let $(\mathcal{X}, \mathcal{D})$ be a model. Then there exist a proper closed subset $Z = Z(X_{\overline{K}}, D_{\overline{K}}) \subset X \setminus D$ and a real constant $m = m(X_{\overline{K}}, D_{\overline{K}}) > 0$ with the following property: for every $\sigma_P \in \mathcal{X}(\mathcal{O}_{S, \mathcal{D}})$ with $\sigma_P(B) \not\subset Z$, where Z denotes the Zariski closure of Z in $\mathcal{X}$, we have

$$\deg_B \sigma_P^* \mathcal{D} \leq m \max\{1, 2g(B) - 2 + \#S\}. \quad (1)$$

Roughly speaking it says that every integral point either lies in a fixed *exceptional* (proper) closed set Z or has height bounded *linearly* in the Euler characteristic $2g(B) - 2 + \#S$, with slope independent of the base.

The geometric Lang-Vojta conjecture has been settled in very few cases: when X is a curve; when $X = \mathbb{P}^2$ and D has a special shape; when $X = A$ is an abelian variety with trivial $K|k$ trace or when $k = \mathbb{C}$ and A is defined over $\mathbb{C}$ (constant case). For a quick review of the known cases we refer the reader to [10, Section 1] and we remark that for a general complex abelian variety — in particular, when $\text{Tr}_{K/\mathbb{C}}(A) \neq 0$ and A is nonconstant — the conjecture remains open.

From now on we fix the base field $k = \mathbb{C}$, a model $(\mathcal{X}, \mathcal{D})$ of (X, D) , and a Riemannian metric ρ on B inducing a path-length ℓ_ρ and a distance function $d_\rho(\cdot, \cdot)$. We shall use two distinct notions of “disk”, and we keep them notationally separate throughout.

- A *(closed topological) disk* in B is a closed subset $\overline{\Delta} \subset B$ homeomorphic to the closed unit disk $\overline{\mathbb{D}} = \{z \in \mathbb{C} : |z| \leq 1\}$, with piecewise- C^1 boundary $\partial \overline{\Delta}$. These are the disks removed from B to form the bordered surface B_0 ; we denote them $\overline{\Delta}_1, \dots, \overline{\Delta}_t$. We do *not* require them to be metric balls—indeed the enlargements performed in §3 produce topological disks that are not metric balls.
- A *metric ball* (or *metric disk*) is a set of the form $\Delta_\rho(x_0, r) := \{b \in B : d_\rho(b, x_0) < r\}$, with closed version $\overline{\Delta}_\rho(x_0, r) := \{b \in B : d_\rho(b, x_0) \leq r\}$, for $x_0 \in B$ and $r > 0$. These appear only in the local integral estimates of §2.2 (e.g. Lemma 2.2) and carry genuine metric content. Any metric ball of sufficiently small radius is one instance of a topological disk.

The notion of $(S, \mathcal{D})$ -integral point can be generalized by allowing S to vary while keeping only its cardinality $s = \#S$ bounded, and by requiring the intersection condition to hold only on the complement of finitely many disjoint closed disks. This notion of generalized integral points was introduced in [10] and we recall it below.

Definition 1.2. Let $t \in \mathbb{Z}_{\geq 0}$. Consider t disjoint disks $\overline{\Delta}_1, \dots, \overline{\Delta}_t$ in B such that $\text{Sing}(f) \subset \bigsqcup_{i=1}^t \overline{\Delta}_i$ and distinct points of $\text{Sing}(f)$ are contained in distinct disks. Set $B_0 := B \setminus \bigsqcup_{i=1}^t \overline{\Delta}_i$, for any $s \in \mathbb{Z}_{\geq 0}$

$$I(s, B_0) := \{\sigma_P \in X(K) : \#(f(\sigma_P(B_0) \cap \mathcal{D})) \leq s\}$$

is the set of *generalized (s, B_0) -integral points* of $(\mathcal{X}, \mathcal{D})$.

One immediately observes the inclusion

$$\bigcup_{\substack{S \subset B \\ \#S \leq s}} \mathcal{X}(\mathcal{O}_{S, \mathcal{D}}) \subseteq I(s, B_0),$$

so $I(s, B_0)$ is potentially much larger than the integral points for any single fixed S . Understanding the growth of $\#I(s, B_0)$ as a function of s (with B_0 fixed) is the main object of this paper.

1.2 Main results

Building on Parshin’s foundational work [9], Phung in [10] proved an important quantitative bound on $\#I(s, B_0)$ for abelian varieties, working in the following setting.

(P) Let A be an abelian variety of dimension n over $K = \mathbb{C}(B)$ and let $\emptyset \neq D \subset A$ be a reduced effective divisor. We fix a model $(\mathcal{A}, \mathcal{D})$, where the proper flat morphism is $f : \mathcal{A} \rightarrow B$ and $\sigma_O : B \rightarrow \mathcal{A}$ is a fixed section. We assume:

- (i) There exist $t \in \mathbb{Z}_{\geq 0}$ disjoint disks $\overline{\Delta}_1, \dots, \overline{\Delta}_t$ in B such that $\text{Sing}(f) \subset \bigsqcup_{i=1}^t \overline{\Delta}_i$ and distinct points of $\text{Sing}(f)$ are contained in distinct disks. We define

$$B_0 := B \setminus \bigsqcup_{i=1}^t \overline{\Delta}_i$$

so that $f: \mathcal{A}_{B_0} \rightarrow B_0$ is a family of abelian varieties $(A_b, \sigma_O(b))_{b \in B_0}$.

- (ii) B_0 is hyperbolic (automatic whenever $t \geq 1$).
 (iii) $D \subset A$ does not contain any translates of nonzero abelian subvarieties.

Phung's main result is the following.

Theorem 1.3 (Phung [10, Theorem A]). *Assume that the setting (P) holds. Then there exists $m := m(\mathcal{A}, \mathcal{D}, B_0) \in \mathbb{R}_{>0}$ such that*

$$\# \left(I(s, B_0) / \text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) \right) \leq m(s+1)^{2n \text{rk}(\pi_1(B_0, b_0))}, \quad \forall s \in \mathbb{Z}_{\geq 0}. \quad (2)$$

We point out that [10, Theorem A] is stated under the additional assumption that D is ample; ampleness, however, is used in [10] only for the height estimate of [10, Corollary 1.3] and plays no role in the proof of Theorem 1.3, which we recall in full in §3.

Note that the bound behaves coherently: if the number of removed disks t increases, then $\text{rk}(\pi_1(B_0, b_0))$ is larger; on the other hand B_0 shrinks, so the set of generalized integral points grows as well.

The proof of Theorem 1.3 rests on a Parshin cocycle argument. Let $k := \text{rk}(\pi_1(B_0, b_0))$ and fix generators $\alpha_1, \dots, \alpha_k$. By Ehresmann's theorem, the fibration $\mathcal{A}_{B_0} \rightarrow B_0$ gives rise to a short exact sequence $1 \rightarrow \Gamma \rightarrow \pi_1(\mathcal{A}_{B_0}, w_0) \rightarrow G \rightarrow 1$, where $\Gamma = H_1(A_{b_0}, \mathbb{Z}) \cong \mathbb{Z}^{2n}$ and $G = \pi_1(B_0, b_0)$. Each section σ_P induces a splitting of this sequence, and the difference between the splittings of σ_P and σ_O defines a 1-cocycle $c_P: G \rightarrow \Gamma$ whose cohomology class determines P modulo the trace and torsion. To count the possible cocycle classes, one bounds the lattice element $c_P(\alpha_j) \in \Gamma \cong \mathbb{Z}^{2n}$ for each generator by the displacement in the universal cover, which in turn is controlled by the h -length of the loop $\sigma_P(\gamma_j)$ in $\mathcal{A}_{B_0}$. A theorem of Green provides the comparison $\ell_h(\sigma_P(\gamma_j)) \leq c^{-1} \ell_S(\gamma_j)$, so the problem reduces to bounding the Kobayashi length of loops γ_j in $B_0 \setminus S$ representing the generators α_j . Phung's linear bound $\ell_S(\gamma_j) = O(s)$ then yields the displacement bound $H(s) = O(s)$, and lattice counting in $\Gamma \cong \mathbb{Z}^{2n}$ gives at most $O(H(s)^{2n}) = O(s^{2n})$ possibilities per generator, hence the exponent $2nk$ in (2).

When D is moreover ample, Theorem 1.3 implies that for every $\sigma_P \in I(s, B_0)$ there exists a constant $M = M(\mathcal{A}, \mathcal{D}, B_0, s) > 0$ such that $\deg_B \sigma_P^* \mathcal{D} < M$ (see [10, Corollary 1.3]). This is a weak form of the geometric Lang–Vojta conjecture for generalized integral points, weak in two compatible senses. On the one hand it holds for the *larger* set $I(s, B_0)$ of generalized integral points, rather than for the integral points attached to a single fixed S . On the other hand the conclusion is correspondingly weaker: Conjecture 1.1 demands a bound that is *linear* in $\#S$ with slope m *independent of s* , whereas $M = M(s)$ is only known to be finite for each fixed s , with a priori no control on its growth as $s \rightarrow \infty$ — the issue is thus the absence of a uniform linear rate, not the mere presence of s . We also stress that [10, Corollary 1.3] assumes D *ample*, which is strictly stronger than the log general type (D big) hypothesis of Conjecture 1.1; as recalled above, ampleness enters only in this corollary and plays no role in Theorem 1.3.

The exponent $2n \text{rk}(\pi_1(B_0))$ in (2) can be compared with known results for the smaller set

$$J(s) := \{ \sigma_P \in A(K) : \#(f(\sigma_P(B) \cap \mathcal{D})) \leq s \} \subseteq I(s, B_0).$$

When $n = 1$ (elliptic curves), classical height-theoretic bounds of Hindry–Silverman [5, Corollary 8.5] together with Shioda's bound [12, Theorem 2.5] for the Mordell–Weil rank show that $\#J(s)$ is bounded by a polynomial in s of degree at most $\text{rk}(\pi_1(B_0))$. This is half the exponent appearing in Phung's bound for the much larger set $I(s, B_0)$.

The main result of this paper eliminates this discrepancy:

Theorem 1.4. *Assume that the setting (P) holds. Then for every $\varepsilon > 0$ there exists $m := m(\mathcal{A}, \mathcal{D}, B_0, \varepsilon) \in \mathbb{R}_{>0}$ such that*

$$\# \left(I(s, B_0) / \text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) \right) \leq m(s+1)^{n \text{rk}(\pi_1(B_0, b_0)) + \varepsilon}, \quad \forall s \in \mathbb{Z}_{\geq 0}. \quad (3)$$

The key ingredient behind [Theorem 1.4](#) is a new, essentially optimal, upper bound for the Kobayashi length of loops on punctured bordered surfaces ([Theorem 1.5](#)). Let us briefly describe the context. Let B_0 be the bordered hyperbolic surface introduced above. For a finite set $S \subset B_0$, the punctured surface $B_0 \setminus S$ carries its own Kobayashi metric, and the induced length ℓ_S of paths in $B_0 \setminus S$ is larger than the length of the same paths measured in the Kobayashi metric of B_0 : the new cusps created by the punctures stretch the metric, so loops become longer as the cardinality of S grows. Given a loop α representing a nontrivial class in $\pi_1(B_0, b_0)$, it is natural to ask how the minimal Kobayashi length of a representative of α in $B_0 \setminus S$ grows with the number of punctures. This is captured by the quantity

$$L(\alpha, s) := \sup_{\substack{S \subset B_0 \\ \#S \leq s}} \inf \{ \ell_S(\gamma) : \gamma \subset B_0 \setminus S \text{ is a loop freely homotopic to } \alpha \text{ in } B_0 \} ,$$

which measures the worst-case minimal Kobayashi length over all configurations of at most s punctures. Phung [10, Theorems B, C] proved that $L(\alpha, s)$ grows at least as $c\sqrt{s}/\log(s+2)$ and at most linearly in s . As explained above, the linear upper bound $L(\alpha, s) = O(s)$ is what produces the factor of 2 in the exponent of (2): replacing $H(s) = O(s)$ by $H(s) = O(\sqrt{s} \log s)$ in the lattice counting reduces the exponent from $2nk$ to $nk + \varepsilon$, for every $\varepsilon > 0$ (the logarithmic factor being absorbed into the ε). We prove:

Theorem 1.5. *Fix a loop $\alpha \subset B_0$ representing a nontrivial class in $\pi_1(B_0, b_0)$. There exists $C > 0$, depending only on α , B_0 , and ρ , such that*

$$L(\alpha, s) \leq C\sqrt{(s+1)\log(s+2)} \quad \text{for all } s \geq 0. \quad (4)$$

Combined with Phung's lower bound, this determines the exact growth rate (Corollary 2.6):

$$\lim_{s \rightarrow +\infty} \frac{\log L(\alpha, s)}{\log s} = \frac{1}{2}. \quad (5)$$

This answers [10, Question 1.4] which asked about the asymptotic behavior of $\frac{\log L(\alpha, s)}{\log s}$.

The proof of [Theorem 1.5](#) proceeds in two steps. First, we establish the bound under the assumption that α admits a smooth *simple* representative $\gamma \in B_0$. We consider the Fermi strip $T_\delta(\gamma)$, a thin tubular neighborhood of γ foliated by parallel curves γ_u (the simplicity of γ ensures that the exponential map from the normal bundle is a diffeomorphism for δ small enough). Phung's linear upper bound $L(\alpha, s) = O(s)$ in [10] is obtained by estimating the Kobayashi length of a single loop directly; our improvement replaces this pointwise approach with an integral averaging argument. The key idea is to bound the L^p -norm of the distortion function λ_S (the supremal directional ratio of the Kobayashi–Royden metric of $B_0 \setminus S$ to the background metric ρ) over the entire strip, for a variable exponent $1 < p < 2$. This is achieved by a Voronoi decomposition of the strip into cells centered at the punctures: near each puncture, λ_S blows up like the inverse of the distance, and the L^p -integrability for $p < 2$ (but not for $p = 2$) controls the singularity. The crucial gain over the linear bound is that each of the s punctures contributes only $O(1/(2-p))$ to the L^p -mass of λ_S on the strip, so the total mass is $O(s/(2-p))$; Hölder's inequality then converts this L^p -bound into an upper bound of order $(s/(2-p))^{1/p}$ —sublinear in s —on the Kobayashi length of a typical parallel curve γ_{u_0} , which is freely homotopic to α in B_0 and avoids S . Optimising the exponent as $p = 2 - 1/\log(s+2)$ balances the divergence of the L^p -norm as $p \rightarrow 2^-$ against the sharpness of the Hölder estimate, yielding the $O(\sqrt{(s+1)\log(s+2)})$ bound.

In the second step, the general case is reduced to the simple case via a word decomposition argument: B_0 has the homotopy type of a compact orientable surface of genus g with t boundary circles, so $\pi_1(B_0, b_0)$ is a free group admitting a basis $\alpha_1, \dots, \alpha_k$ of *smooth simple loops* (the generators around the disks, together with the standard genus generators when $g \geq 1$); any nontrivial $\alpha \in \pi_1(B_0, b_0)$ can be written as a word $\alpha_{i_1}^{\varepsilon_1} \dots \alpha_{i_m}^{\varepsilon_m}$ in this basis. Applying the simple case to each basis loop and concatenating the resulting parallel curves at a common base point (which can be arranged, see Proposition 2.5 below) yields a representative of α avoiding S whose Kobayashi length is at most m times the bound for a single basis loop.

[Theorem 1.4](#) follows from [Theorem 1.5](#) by substituting the improved length estimate into the Parshin cocycle argument outlined above. The cocycle construction requires loops representing the generators $\alpha_1, \dots, \alpha_k$ that share a *common base point* and are conjugated to the generators by a *single* path. [Theorem 1.5](#), applied independently to each generator, produces loops with the optimal Kobayashi length bound but with potentially different base points. In Proposition 2.5 we show that, by choosing smooth simple representatives of $\alpha_1, \dots, \alpha_k$ that share a common tangent direction at b_0 (so that their Fermi

strips can be controlled by a single averaging parameter), a common base point can be arranged without degrading the asymptotic bound. The improved displacement bound $H(s) = O(\sqrt{s} \log s)$ then propagates through the lattice counting argument unchanged, halving the exponent.

Funding. The author is supported by the NSFC RFIS-I 2025, project titled “*Interdisciplinary Perspectives in Diophantine Geometry: Analytic, Dynamical and Arithmetic Explorations*”, and by the Zhejiang Science and Technology Development Fund [2024] No. 28, project titled “*New Perspectives in Diophantine Geometry*”.

2 Kobayashi lengths estimates

2.1 Kobayashi pseudo-metric

Let X be a connected complex manifold and let TX be its tangent bundle. Throughout, TX denotes the *real* tangent bundle $T^{\mathbb{R}}X$, regarded as a complex vector bundle (of complex rank $\dim_{\mathbb{C}} X$) via the complex structure J , so that $\lambda v := (\operatorname{Re} \lambda) v + (\operatorname{Im} \lambda) Jv$ for $\lambda \in \mathbb{C}$ and $v \in T_{X,x}^{\mathbb{R}}$; this is canonically $\mathbb{C}$ -isomorphic to the holomorphic tangent bundle $T^{1,0}X$ and we use the two interchangeably. In particular it is *not* the complexification $T^{\mathbb{R}}X \otimes_{\mathbb{R}} \mathbb{C} = T^{1,0}X \oplus T^{0,1}X$, whose complex rank is twice as large.

Definition 2.1. A *Finsler pseudo-metric* on X is a function

$$\begin{aligned} F: TX &\rightarrow \mathbb{R}_{\geq 0} \\ (x, v) &\mapsto F(x, v) \end{aligned}$$

satisfying the homogeneity condition $F(x, \lambda v) = |\lambda| F(x, v)$ for all $\lambda \in \mathbb{C}$. If $F(x, v) > 0$ for all $x \in X$ and $v \in T_{X,x} \setminus \{0\}$, then we say that F is a *Finsler metric*.

Throughout, by a *path* we mean a piecewise C^1 map $\gamma: [0, 1] \rightarrow X$; this entails no loss of generality for the distances defined below. Given a Borel measurable Finsler pseudo-metric F and a path γ , the function $t \mapsto F(\gamma(t), \gamma'(t))$ is measurable, and we define the length of γ as

$$\ell_F(\gamma) := \int_0^1 F(\gamma(t), \gamma'(t)) dt \in [0, +\infty].$$

When F is *upper semicontinuous* we have $\ell_F(\gamma) < +\infty$, since F is then locally bounded above and $\gamma([0, 1])$ is compact. Given a set of paths Γ we set

$$\ell_F(\Gamma) := \inf_{\gamma \in \Gamma} \ell_F(\gamma).$$

We endow X with the structure of extended pseudo-metric space by setting

$$d_F(x, y) := \inf_{\gamma} \ell_F(\gamma), \quad \text{where the infimum runs over all paths } \gamma \text{ joining } x \text{ and } y.$$

Moreover, if Y, Z are two subsets of X , we put

$$\operatorname{diam}_F(Y) := \sup_{y, y' \in Y} d_F(y, y'), \quad d_F(Y, Z) := \inf_{y \in Y, z \in Z} d_F(y, z).$$

One can show that a *continuous* Finsler metric induces a metric d_F . Any Hermitian metric h on X induces a Finsler metric $F_h(x, v) = \sqrt{h_x(v, v)}$, hence a distance which we denote by d_h .

An important example of a Finsler pseudo-metric is the *Kobayashi–Royden pseudo-metric*

$$\kappa_X(x, v) := \inf \left\{ \frac{2}{R} \in \mathbb{R}_{>0} : \exists f \in \operatorname{Hol}(\Delta(0, R), X) \text{ such that } f(0) = x, f'(0) = v \right\},$$

where $\operatorname{Hol}(\Delta(0, R), X)$ is the set of holomorphic maps from the open disk of radius R centered at 0 to X . Royden [11, Proposition 3] shows that κ_X is upper semicontinuous, hence Borel measurable. Since κ_X is intrinsic to X , the induced topological pseudo-metric is denoted by d_X and called the *Kobayashi pseudo-metric*. We say that X is *Kobayashi hyperbolic* if d_X is a metric.

If X carries a Hermitian metric h whose holomorphic sectional curvature is bounded above by a negative constant, then by the Ahlfors–Schwarz lemma one has $\kappa_X \geq c \|\cdot\|_h$ for some constant $c > 0$; in

particular $d_X \geq c d_h$ and X is Kobayashi hyperbolic. In dimension ≥ 2 the converse is a delicate matter and is not known to hold in general. On a Riemann surface, however, it holds: if X is Kobayashi hyperbolic its universal cover is the disk, so X is biholomorphic to $\Gamma \backslash \mathbb{H}$ for a torsion-free Fuchsian group $\Gamma \cong \pi_1(X)$ (no elliptic elements, parabolics allowed). The curvature -1 Poincaré metric $ds^2 = (dx^2 + dy^2)/y^2$ on $\mathbb{H}$ descends to X , and since holomorphic covering maps are infinitesimal isometries for κ_X , the descended metric *is* the Kobayashi metric. Thus on Riemann surfaces “admits a Hermitian metric of holomorphic sectional curvature bounded above by a negative constant” and “Kobayashi hyperbolic” coincide.

From the definition one deduces the distance-decreasing property of the Kobayashi–Royden metric: if $\phi: X \rightarrow Y$ is holomorphic, then

$$\kappa_Y(\phi(x), d_x \phi(v)) \leq \kappa_X(x, v), \quad \forall (x, v) \in TX.$$

This in turn implies the distance-decreasing property of the Kobayashi distance

$$d_Y(\phi(x), \phi(x')) \leq d_X(x, x'), \quad \forall x, x' \in X.$$

2.2 Growth of loop lengths in terms of the cusps

We fix a compact Riemann surface B of genus g endowed with a Riemannian metric ρ . Let $\overline{\Delta}_1, \dots, \overline{\Delta}_t$ be disjoint disks on B and define $B_0 := B \setminus \sqcup_{i=1}^t \overline{\Delta}_i$. We assume that $t \geq 1$, so that B_0 is hyperbolic. Let $S \subseteq B_0$ be a finite set (possibly empty), put $s := \#S$, and let κ_S be the Kobayashi–Royden metric of the hyperbolic surface $B_0 \setminus S$, with induced length and distance ℓ_S, d_S . We define the *distortion function*

$$\lambda_S(b) := \sup_{v \in T_b B \setminus \{0\}} \frac{\kappa_S(b, v)}{\sqrt{\rho_b(v, v)}}, \quad b \in B_0 \setminus S, \quad (6)$$

where $T_b B = T_b^{\mathbb{R}} B$ is the *real* tangent plane (real dimension 2), on which both $\kappa_S(b, \cdot)$ and $\sqrt{\rho_b(\cdot, \cdot)}$ are defined. The ratio is homogeneous of degree 0, so the supremum may be taken over the compact ρ -unit circle, where it is the upper semicontinuous function $\kappa_S(b, \cdot)$; thus $\lambda_S(b)$ is finite and attained, and λ_S is Borel on $B_0 \setminus S$. In a holomorphic coordinate $z = x + iy$, identifying $a\partial_x + b'\partial_y \leftrightarrow (a + ib')\partial_z$, $\mathbb{C}$ -homogeneity gives $\kappa_S(b, a\partial_x + b'\partial_y) = \kappa_S(b, \partial_z) \sqrt{a^2 + b'^2}$, while ρ_b has eigenvalues $\mu_1(b) \leq \mu_2(b)$ in the frame (∂_x, ∂_y) . Maximizing the ratio over the Euclidean unit circle thus minimizes $\rho_b(v, v)$, giving

$$\lambda_S(b) = \frac{\kappa_S(b, \partial_z)}{\sqrt{\mu_1(b)}}.$$

If ρ is conformal ($\mu_1 = \mu_2$, or equivalently ρ is hermitian) the ratio of Equation (6) is independent of v (so that the supremum on v is superfluous) and λ_S is smooth; we do not assume this.

Standing assumption for §2.2 (lifted in §2.3). Throughout this subsection, we fix a loop α in B_0 that doesn't represent a trivial class in $\pi_1(B_0, b_0)$ and that admits a smooth simple representative $\gamma \in B_0$ freely homotopic to α . Here and throughout, a *smooth simple loop* is an injective smooth immersion $\gamma: S^1 \rightarrow B_0$; since S^1 is compact, γ is then a smooth embedding, and its image is a smoothly embedded circle. In particular γ' never vanishes, so γ may be parametrized by ρ -arc length, with $\gamma'(0) = \gamma'(\ell_\rho(\gamma))$. Such loops exist: the basis loops of $\pi_1(B_0, b_0)$ described in §2.3 are smooth simple, and the general case (no simplicity hypothesis) is reduced to this one in §2.3.

Parametrize γ by its ρ -arc-length and let $\eta(\tau)$ be the unit normal vector of γ at τ ; here and throughout we fix the convention that $\eta(\tau)$ is the rotation of the unit tangent vector $\gamma'(\tau)$ by $+\pi/2$, with respect to ρ and the orientation of the Riemann surface B . Since γ is smooth and simple and compactly contained in B_0 , for $\delta \in \mathbb{R}_{>0}$ sufficiently small the exponential map from the normal bundle is a diffeomorphism, yielding:

$$\begin{aligned} \mathbb{R}/\ell_\rho(\gamma)\mathbb{Z} \times [-\delta, \delta] &\rightarrow \{b \in B_0 : d_\rho(b, \gamma) \leq \delta\} =: T_\delta(\gamma) \subseteq B_0 \\ (\tau, u) &\mapsto \exp_{\gamma(\tau)}(u\eta(\tau)) \end{aligned}$$

The coordinates (τ, u) are called *Fermi coordinates*, and $T_\delta(\gamma)$ is the *Fermi strip* with respect to γ . For any $u \in [-\delta, \delta]$ we define the *parallel curve* $\gamma_u: [0, \ell_\rho(\gamma)] \rightarrow B_0$, $\gamma_u(\tau) := \exp_{\gamma(\tau)}(u\eta(\tau))$.

In Fermi coordinates the induced metric on each parallel curve γ_u is $\rho|_{\gamma_u} = \rho_{\tau\tau} d\tau^2$, so its ρ -length is

$$\ell_\rho(\gamma_u) = \int_0^{\ell_\rho(\gamma)} \sqrt{\rho_{\tau\tau}(\tau, u)} d\tau.$$




Figure 1: The Fermi strip $T_\delta(\gamma)$ around γ , with punctures $p_1, \dots, p_5 \in S$ and the parallel curve γ_u .

Write the Riemannian area form in Fermi coordinates as $dA_\rho = J(\tau, u) d\tau du$, where

$$J(\tau, u) := \sqrt{\rho_{\tau\tau}\rho_{uu} - \rho_{\tau u}^2}$$

is the square root of the determinant of the metric ρ in the coordinates (τ, u) . We claim that, for δ small enough, J and $\sqrt{\rho_{\tau\tau}}$ are uniformly comparable on the strip. Indeed, by the properties of Fermi coordinates ([6, Proposition 5.26]), the metric at $u = 0$ satisfies $\rho_{\tau u}(\tau, 0) = 0$ and $\rho_{uu}(\tau, 0) = 1$, so $J(\tau, 0) = \sqrt{\rho_{\tau\tau}(\tau, 0)}$. Now $\rho_{\tau\tau}(\cdot, 0)$ is continuous and attains a positive minimum on the compact circle $[0, \ell_\rho(\gamma)] \times \{0\}$; by uniform continuity of $\rho_{\tau\tau}$ there is a $\delta > 0$ —which we also shrink, if necessary, so that the exponential map of the normal bundle is a diffeomorphism onto $T_\delta(\gamma)$ —such that $\rho_{\tau\tau} > 0$ on the whole strip $[0, \ell_\rho(\gamma)] \times [-\delta, \delta]$. Then J and $\sqrt{\rho_{\tau\tau}}$ are continuous and strictly positive there, so the ratio $J/\sqrt{\rho_{\tau\tau}}$ is a continuous positive function on a compact set. It follows that there exists $\kappa_0 = \kappa_0(\rho, \gamma, \delta) \geq 1$ such that

$$\kappa_0^{-1} \sqrt{\rho_{\tau\tau}(\tau, u)} \leq J(\tau, u) \leq \kappa_0 \sqrt{\rho_{\tau\tau}(\tau, u)} \quad \forall (\tau, u) \in [0, \ell_\rho(\gamma)] \times [-\delta, \delta]. \quad (7)$$

In particular, for any measurable $f \geq 0$:

$$\kappa_0^{-1} \int_{-\delta}^{\delta} \int_0^{\ell_\rho(\gamma)} f(\tau, u) \sqrt{\rho_{\tau\tau}(\tau, u)} d\tau du \leq \int_{T_\delta(\gamma)} f dA_\rho \leq \kappa_0 \int_{-\delta}^{\delta} \int_0^{\ell_\rho(\gamma)} f(\tau, u) \sqrt{\rho_{\tau\tau}(\tau, u)} d\tau du. \quad (8)$$

Lemma 2.2. *There exists a constant $A := A(\rho, B_0, \alpha) \in \mathbb{R}_{>0}$ such that for every $1 \leq p < 2$ and every finite set $S \subset B_0$:*

$$\int_{T_\delta(\gamma)} \lambda_S^p dA_\rho \leq \frac{A(s+1)}{2-p}.$$

Proof. Throughout this proof, $\Delta(\cdot, \cdot) = \Delta_\rho(\cdot, \cdot)$ denotes a metric ball for the Riemannian distance d_ρ (not a removed topological disk of B_0). Assume $S = \{p_1, \dots, p_s\} \neq \emptyset$. Consider the Voronoi decomposition of $T_\delta(\gamma)$ induced by S : for each $p_j \in S$, define

$$V_j := \{b \in T_\delta(\gamma) : d_\rho(b, p_j) \leq d_\rho(b, p_i) \forall i \neq j\}.$$

Note that the punctures p_j need not lie inside $T_\delta(\gamma)$. Nevertheless, the cells cover the strip: $T_\delta(\gamma) = \bigcup_{j=1}^s V_j$ (some cells V_j may be empty, and distinct cells may overlap along tie sets).

Step 1: Pointwise and area bounds. Put $T := T_\delta(\gamma)$ and fix $b \in T \setminus S$. Set $D_b := d_\rho(b, S) > 0$. By [10, Lemma 3.3], there exist constants $c_1, r_1 > 0$ (depending only on B and ρ) such that for every $b \in B$, every $v \in T_b B \setminus \{0\}$, and every $0 < r < r_1$:

$$\kappa_{\Delta(b,r)}(b, v) \leq \frac{c_1}{r} \sqrt{\rho_b(v, v)}.$$

By [10, Lemma 3.5], there exist constants $c_2, r_2 > 0$ such that $\text{Area}_\rho(\Delta(q, r)) \leq c_2 r^2$ for all $q \in B$ and $0 < r \leq r_2$. Set $D := \min\{d_\rho(T, \partial B_0), r_1, r_2\} > 0$ and $R_b := \min\{D_b, D\}$. Then $\Delta(b, R_b) \subset B_0 \setminus S$ (it lies in B_0 because $R_b \leq D \leq d_\rho(T, \partial B_0)$, and avoids S because $R_b \leq D_b$), so the decreasing property of the Kobayashi–Royden metric (applied to the inclusion $\Delta(b, R_b) \hookrightarrow B_0 \setminus S$) gives, for every $v \in T_b B \setminus \{0\}$, the bound $\kappa_{B_0 \setminus S}(b, v) \leq \kappa_{\Delta(b, R_b)}(b, v)$. Taking the supremum over v and using that [10, Lemma 3.3] holds uniformly in v :

$$\lambda_S(b) = \sup_{v \neq 0} \frac{\kappa_{B_0 \setminus S}(b, v)}{\sqrt{\rho_b(v, v)}} \leq \sup_{v \neq 0} \frac{\kappa_{\Delta(b, R_b)}(b, v)}{\sqrt{\rho_b(v, v)}} \leq \frac{c_1}{R_b} = \frac{c_1}{\min\{D_b, D\}}. \quad (9)$$

Moreover, since $D \leq r_2$:

$$\text{Area}_\rho(\Delta(q, r)) \leq c_2 r^2 \quad \text{for all } q \in B \text{ and } r \leq D. \quad (10)$$

Step 2: Integral over Voronoi cells. For $b \in V_j$, the point p_j is the nearest element of S to b , so $D_b = d_\rho(b, p_j)$. We split V_j into two regions:

$$V_j^{\text{near}} := \{b \in V_j : d_\rho(b, p_j) \leq D\}, \quad V_j^{\text{far}} := \{b \in V_j : d_\rho(b, p_j) > D\}.$$

On V_j^{near} we have $R_b = D_b = d_\rho(b, p_j)$, so (9) gives $\lambda_S(b) \leq c_1/d_\rho(b, p_j)$. By the “layer cake representation” in measure theory ($\int f^p d\mu = p \int_0^\infty t^{p-1} \mu(\{f > t\}) dt$, see [7, Theorem 1.13]) and the substitution $r = 1/t$:

$$\begin{aligned} \int_{V_j^{\text{near}}} \lambda_S^p dA_\rho &\leq c_1^p \int_{V_j^{\text{near}}} \frac{dA_\rho}{d_\rho(b, p_j)^p} \\ &= p c_1^p \int_0^{+\infty} t^{p-1} \text{Area}_\rho\{b \in V_j^{\text{near}} : d_\rho(b, p_j)^{-1} > t\} dt \\ &= p c_1^p \int_0^{+\infty} \frac{\text{Area}_\rho(V_j^{\text{near}} \cap \Delta(p_j, r))}{r^{p+1}} dr \\ &\leq p c_1^p \int_0^D \frac{\text{Area}_\rho(\Delta(p_j, r))}{r^{p+1}} dr + p c_1^p \int_D^{+\infty} \frac{\text{Area}_\rho(T)}{r^{p+1}} dr \\ &\leq \frac{p c_1^p c_2 D^{2-p}}{2-p} + \frac{c_1^p \text{Area}_\rho(T)}{D^p}, \end{aligned} \tag{11}$$

where the fourth line uses $V_j^{\text{near}} \cap \Delta(p_j, r) \subset \Delta(p_j, r)$ for $r \leq D$, and $\text{Area}_\rho(V_j^{\text{near}} \cap \Delta(p_j, r)) \leq \text{Area}_\rho(T)$ for $r \geq D$. Moreover the fifth line uses the area bound (10) and $\int_D^{+\infty} r^{-(p+1)} dr = D^{-p}/p$. Note that the integral $\int_0^D r^{1-p} dr$ converges since $p < 2$.

On V_j^{far} we have $R_b = D$, so $\lambda_S(b) \leq c_1/D$ and

$$\int_{V_j^{\text{far}}} \lambda_S^p dA_\rho \leq \frac{c_1^p}{D^p} \text{Area}_\rho(T). \tag{12}$$

Set $c_3 := \max\{2 \max\{1, c_1^2\} c_2 \max\{1, D\}, 2 \max\{c_1/D, c_1^2/D^2\} \text{Area}_\rho(T)\}$. Since $1 \leq p < 2$ we have $c_1^p \leq \max\{1, c_1^2\}$, $D^{2-p} \leq \max\{1, D\}$, and $c_1^p/D^p \leq \max\{c_1/D, c_1^2/D^2\}$, so combining (11) and (12):

$$\int_{V_j} \lambda_S^p dA_\rho \leq \frac{c_3}{2-p} + c_3 \leq \frac{2c_3}{2-p}, \tag{13}$$

where the last inequality uses $1 \leq 1/(2-p)$ for $p \geq 1$.

Step 3: Summing over cells. Since the cells V_j cover T and $\lambda_S^p \geq 0$, subadditivity of the integral gives (cells with $V_j = \emptyset$ contribute zero):

$$\int_T \lambda_S^p dA_\rho \leq \sum_{j=1}^s \int_{V_j} \lambda_S^p dA_\rho \leq \frac{2c_3 s}{2-p}.$$

If $s = 0$ (i.e. $S = \emptyset$) on T we have

$$\lambda_S(b) = \sup_{v \neq 0} \frac{\kappa_{B_0}(b, v)}{\sqrt{\rho_b(v, v)}} \leq M := \sup_{b \in T} \sup_{v \neq 0} \frac{\kappa_{B_0}(b, v)}{\sqrt{\rho_b(v, v)}} < +\infty;$$

where the finiteness follows from $T \subseteq B_0$. In this case we have

$$\int_T \lambda_S^p dA_\rho \leq M^p \text{Area}_\rho(T) \leq \max\{1, M^2\} \text{Area}_\rho(T).$$

Setting $A := \max\{2c_3, \max\{1, M^2\} \text{Area}_\rho(T)\}$ the claim is proved. $\square$

Let us now introduce the main object of this section and then prove the main result.

Definition 2.3. Fix a loop α in B_0 that doesn't represent a trivial class in $\pi_1(B_0, b_0)$. Let Γ_S^α denote the set of loops in $B_0 \setminus S$ that are freely homotopic to α in B_0 . We define

$$L(\alpha, s) := \sup_{\substack{S \subseteq B_0 \\ \#S \leq s}} \ell_S(\Gamma_S^\alpha).$$

Note that since α is chosen (non-trivially) in B_0 and the free homotopies are checked in the non-cuspidal hyperbolic surface B_0 , we have $\ell_S(\Gamma_S^\alpha) > 0$ by [?, Proposition 2.24].

Proof of Theorem 1.5 under the standing simplicity assumption. Consider the smooth simple loop $\gamma \in B_0$ freely homotopic to α , fixed by the standing assumption at the beginning of the section. Put $T := T_\delta(\gamma)$ parametrized with Fermi coordinates (τ, u) as above. Set

$$L_0 := \sup_{|u| \leq \delta} \ell_\rho(\gamma_u) < +\infty.$$

Fix an arbitrary $S \subset B_0$ with $\#S = s$ and a parameter $1 < p < 2$. Define $q := \frac{p}{p-1}$ (the conjugate exponent, so $1/p + 1/q = 1$). Recall that in Fermi coordinates $\gamma'_u(\tau) = \partial_\tau|_{(\tau, u)}$ and $\rho_{\tau\tau}(\tau, u) = \rho_{\gamma_u(\tau)}(\gamma'_u(\tau), \gamma'_u(\tau))$, so by the definition of λ_S as a directional supremum,

$$\kappa_S(\gamma_u(\tau), \gamma'_u(\tau)) \leq \lambda_S(\gamma_u(\tau)) \sqrt{\rho_{\gamma_u(\tau)}(\gamma'_u(\tau), \gamma'_u(\tau))} = \lambda_S(\tau, u) \sqrt{\rho_{\tau\tau}(\tau, u)}.$$

For each $u \in]-\delta, \delta[$ with $\gamma_u \cap S = \emptyset$, Hölder's inequality with respect to the measure $d\mu = \sqrt{\rho_{\tau\tau}} d\tau$ then gives:

$$\begin{aligned} \ell_S(\gamma_u) &= \int_0^{\ell_\rho(\gamma)} \kappa_S(\gamma_u(\tau), \gamma'_u(\tau)) d\tau \\ &\leq \int_0^{\ell_\rho(\gamma)} \lambda_S(\tau, u) \sqrt{\rho_{\tau\tau}(\tau, u)} d\tau \\ &\leq \left(\int_0^{\ell_\rho(\gamma)} \lambda_S(\tau, u)^p \sqrt{\rho_{\tau\tau}} d\tau \right)^{1/p} \left(\int_0^{\ell_\rho(\gamma)} \sqrt{\rho_{\tau\tau}} d\tau \right)^{1/q} \\ &= \left(\int_0^{\ell_\rho(\gamma)} \lambda_S(\tau, u)^p \sqrt{\rho_{\tau\tau}} d\tau \right)^{1/p} \ell_\rho(\gamma_u)^{1/q}. \end{aligned} \quad (14)$$

By Equation (8) and Lemma 2.2 we get:

$$\int_{-\delta}^{\delta} \int_0^{\ell_\rho(\gamma)} \lambda_S(\tau, u)^p \sqrt{\rho_{\tau\tau}} d\tau du \leq \kappa_0 \int_T \lambda_S^p dA_\rho \leq \frac{\kappa_0 A(s+1)}{2-p}. \quad (15)$$

The set $E := \{u \in]-\delta, \delta[: \gamma_u \cap S \neq \emptyset\}$ has at most s points, hence has measure zero. Define $g(u) := \int_0^{\ell_\rho(\gamma)} \lambda_S(\tau, u)^p \sqrt{\rho_{\tau\tau}} d\tau$ and $M := \frac{\kappa_0 A(s+1)}{2\delta(2-p)}$. By (15),

$$\frac{1}{2\delta} \int_{-\delta}^{\delta} g(u) du \leq M. \quad (16)$$

We claim there exists $u_0 \in]-\delta, \delta[\setminus E$ with $g(u_0) \leq M$. If not, then $g(u) > M$ for all $u \in]-\delta, \delta[\setminus E$. Since $g \geq 0$ and E has measure zero:

$$\frac{1}{2\delta} \int_{-\delta}^{\delta} g(u) du = \frac{1}{2\delta} \int_{]-\delta, \delta[\setminus E} g(u) du > \frac{1}{2\delta} \cdot M \cdot 2\delta = M,$$

contradicting (16). Hence such u_0 exists, and since $u_0 \notin E$ we have $\gamma_{u_0} \cap S = \emptyset$ and

$$\int_0^{\ell_\rho(\gamma)} \lambda_S(\tau, u_0)^p \sqrt{\rho_{\tau\tau}(\tau, u_0)} d\tau = g(u_0) \leq M = \frac{\kappa_0 A(s+1)}{2\delta(2-p)}. \quad (17)$$

Substituting Equation (17) into Equation (14) and using $\ell_\rho(\gamma_{u_0}) \leq L_0$ we get

$$\ell_S(\gamma_{u_0}) \leq \left(\frac{\kappa_0 A}{2\delta} \right)^{1/p} L_0^{1/q} \left(\frac{s+1}{2-p} \right)^{1/p}. \quad (18)$$

Since the Fermi coordinate map is a diffeomorphism, the family $\{\gamma_u\}_{u \in [0, u_0]}$ is a free homotopy in $T_\delta(\gamma) \subset B_0$ from $\gamma_0 = \gamma$ to γ_{u_0} . As γ is freely homotopic to α in B_0 (by construction), γ_{u_0} is freely homotopic to α in B_0 by transitivity. Moreover $\gamma_{u_0} \subset B_0 \setminus S$ (since $u_0 \notin E$), so $\gamma_{u_0} \in \Gamma_S^\alpha$.

We now assume $s \geq 1$ (the case $s = 0$ is handled below) and set

$$p := 2 - \frac{1}{\log(s+2)}, \quad \text{so that} \quad 2-p = \frac{1}{\log(s+2)}, \quad p > 1.$$

Then

$$\frac{s+1}{2-p} = (s+1) \log(s+2), \quad \frac{1}{p} = \frac{\log(s+2)}{2 \log(s+2) - 1}.$$

We split

$$\left(\frac{s+1}{2-p}\right)^{1/p} = ((s+1) \log(s+2))^{1/2} \cdot ((s+1) \log(s+2))^{1/p-1/2}$$

and claim the second factor is $O(1)$. Indeed, $1/p - 1/2 = O(1/\log s)$ and

$$\log((s+1) \log(s+2)) = \log(s+1) + \log \log(s+2) = O(\log s),$$

so

$$((s+1) \log(s+2))^{1/p-1/2} = \exp(O(1/\log s) \cdot O(\log s)) = \exp(O(1)) = O(1).$$

Consider now the first factor of Equation (18), namely $C_0(p) := (\kappa_0 A / (2\delta))^{1/p} L_0^{1/q}$. As $p \rightarrow 2^-$ we have $1/p \rightarrow 1/2$ and $1/q \rightarrow 1/2$, so $C_0(p) \rightarrow (\kappa_0 A / (2\delta))^{1/2} L_0^{1/2}$. Since C_0 is continuous on $[2 - 1/\log 3, 2[$ and has a finite limit at $p = 2$, it is bounded there.

Combining, we conclude that, *under the standing assumption that α admits a smooth simple representative $\gamma \in B_0$* , there exists $C > 0$, depending only on α , B_0 , and ρ , such that for every $S \subset B_0$ with $\#S = s \geq 1$:

$$\ell_S(\Gamma_S^\alpha) \leq \ell_S(\gamma_{u_0}) \leq C \sqrt{(s+1) \log(s+2)}.$$

Since this holds for every such S , and since sets with $\#S = s' < s$ satisfy the same bound a fortiori ($s \mapsto \sqrt{(s+1) \log(s+2)}$ being increasing), taking the supremum over all $S \subset B_0$ with $\#S \leq s$ gives $L(\alpha, s) \leq C \sqrt{(s+1) \log(s+2)}$. For $s = 0$, $L(\alpha, 0) = \inf_{\beta \in B_0} \ell_{\kappa_{B_0}}(\beta)$ is a fixed positive constant, which is absorbed into C . $\square$

For the application to generalized integral points, the Parshin cocycle argument requires loops representing the generators of $\pi_1(B_0, b_0)$ that share a *common base point* and are conjugated to the generators by a *single* path. Theorem 1.5, applied independently to each generator, produces loops with the optimal Kobayashi length bound but with potentially different base points. The next proposition shows that a common base point can be arranged without degrading the asymptotic bound.

Lemma 2.4. *Let M be a smooth orientable Riemannian surface without boundary, let $p \in M$, and let $w \in T_p M$ be a unit tangent vector. Then for every smooth simple loop γ at p with $\gamma \in M$, there exists a smooth simple loop $\tilde{\gamma}$ at p with $\tilde{\gamma} \in M$, homotopic to γ relative to p , and satisfying*

$$\frac{\tilde{\gamma}'(0)}{|\tilde{\gamma}'(0)|} = w.$$

Proof. Let $w_0 := \gamma'(0)/|\gamma'(0)|$ be the unit tangent vector of γ at p . If $w_0 = w$ there is nothing to prove, so assume $w_0 \neq w$. Let $\theta \in]0, 2\pi[$ be the angle from w_0 to w and for each $t \in [0, 1]$ let $R_t \in \text{SO}(2)$ denote the rotation of $T_p M$ by angle $t\theta$, so that $R_0 = \text{Id}$ and $R_1(w_0) = w$.

Choose $\epsilon > 0$ small enough that the exponential map $\exp_p$ is a diffeomorphism from $B(0, \epsilon) \subset T_p M$ onto an open neighborhood U of p in M with $U \Subset M$. Choose a smooth cutoff function $\chi: [0, \infty) \rightarrow [0, 1]$ with $\chi \equiv 1$ on $[0, \epsilon/3]$ and $\chi \equiv 0$ on $[2\epsilon/3, \infty)$. Define the map $\phi: M \rightarrow M$ by

$$\phi(x) := \begin{cases} \exp_p(R_{\chi(|y|)} y), & \text{if } x = \exp_p(y) \in U, \\ x, & \text{if } x \notin U, \end{cases}$$

where $|y|$ denotes the norm in $T_p M$. Since $\chi \equiv 1$ near $|y| = 0$, the map ϕ equals $\exp_p \circ R_1 \circ \exp_p^{-1}$ in a neighborhood of p , and since $\chi \equiv 0$ for $|y| \geq 2\epsilon/3$, ϕ is the identity outside U . In geodesic polar coordinates (r, ψ) centered at p (via $\exp_p$), the map reads $\phi(r, \psi) = (r, \psi + \chi(r)\theta)$: it preserves the radius r and rotates each geodesic circle $\{r = \text{const}\}$ by the angle $\chi(r)\theta$. On each circle this is a bijection, and distinct circles map to distinct circles, so ϕ is a bijection of U fixing p ; together with $\phi = \text{id}$ outside U this makes ϕ a bijection of M . It is smooth with smooth inverse $\phi^{-1}(r, \psi) = (r, \psi - \chi(r)\theta)$ (both expressions are smooth across $r = 0$ because $\chi \equiv 1$ there, where ϕ is the fixed rotation R_1), so ϕ is a diffeomorphism of M . Moreover:

- (i) $\phi(p) = p$ and $d_p \phi = R_1$, so $(\phi \circ \gamma)'(0) = R_1(\gamma'(0)) = |\gamma'(0)| w$, which has unit tangent direction w ;

- (ii) ϕ is the identity outside U , so $\phi \circ \gamma$ coincides with γ outside U and in particular $\phi \circ \gamma \in M$;
- (iii) ϕ is a diffeomorphism, so $\phi \circ \gamma$ is simple;
- (iv) the family ϕ_t (defined by replacing χ with $t\chi$ for $t \in [0, 1]$) satisfies $\phi_0 = \text{Id}$ and $\phi_1 = \phi$, and $\phi_t(p) = p$ for all t . Hence $t \mapsto \phi_t \circ \gamma$ is a homotopy of loops based at p from γ to $\tilde{\gamma}$, so $[\tilde{\gamma}] = [\gamma]$ in $\pi_1(M, p)$.

Setting $\tilde{\gamma} := \phi \circ \gamma$ completes the proof. $\square$

Proposition 2.5. *Let B_0 be as above and fix a simple basis $\alpha_1, \dots, \alpha_k$ of $\pi_1(B_0, b_0)$. There exist constants $C, \delta' > 0$, depending only on $\alpha_1, \dots, \alpha_k$, B_0 , and ρ , such that for every finite set $S \subset B_0$ with $\#S = s \geq 1$, there exist a point $q \in B_0 \setminus S$, a path σ from b_0 to q in B_0 with $\ell_\rho(\sigma) \leq \delta'$, and loops $\hat{\gamma}_1, \dots, \hat{\gamma}_k \subset B_0 \setminus S$ based at q such that*

$$[\sigma^{-1} \circ \hat{\gamma}_j \circ \sigma] = \alpha_j \quad \text{in } \pi_1(B_0, b_0) \quad \text{for every } j = 1, \dots, k,$$

and

$$\ell_S(\hat{\gamma}_j) \leq C \sqrt{(s+1) \log(s+2)} \quad \text{for all } j = 1, \dots, k.$$

Moreover, σ and the loops $\hat{\gamma}_1, \dots, \hat{\gamma}_k$ are contained in a fixed compact set $K_0 \Subset B_0$ and satisfy $\ell_\rho(\hat{\gamma}_j) \leq L_0$, where K_0 and $L_0 > 0$ depend only on $\alpha_1, \dots, \alpha_k$, B_0 , and ρ (not on S).

Proof. Fix a unit tangent vector $w \in T_{b_0} B_0$ and let η_0 be the rotation of w by $+\pi/2$ with respect to ρ and the orientation of B . By Lemma 2.4, for each $j = 1, \dots, k$ we can choose a smooth simple loop $\gamma^{(j)}$ at b_0 , compactly contained in B_0 , homotopic to α_j relative to b_0 , and whose unit tangent vector at b_0 equals w . Since all loops share the tangent direction w at b_0 , and since the unit normal field along every loop is the $+\pi/2$ -rotation of the unit tangent (the convention fixed in §2.2), their unit normal vectors at b_0 all equal η_0 . For each j , construct the Fermi strip $T_\delta(\gamma^{(j)})$ as in the proof of Theorem 1.5, with a common width $\delta > 0$ small enough for all k strips. The parallel curve $\gamma_u^{(j)}$ has base point

$$q(u) := \exp_{b_0}(u \eta_0),$$

which is independent of j . For $1 < p < 2$, define

$$g_j(u) := \int_0^{\ell_\rho(\gamma^{(j)})} \lambda_S(\tau, u)^p \sqrt{\rho_{\tau\tau}} d\tau$$

and set $g := \sum_{j=1}^k g_j$. Let $E_j := \{u \in]-\delta, \delta[: \gamma_u^{(j)} \cap S \neq \emptyset\}$; then $\#E_j \leq s$, so the set $E := \bigcup_{j=1}^k E_j$ has at most ks points and in particular has measure zero. By (8) and Lemma 2.2 applied to each Fermi strip,

$$\frac{1}{2\delta} \int_{-\delta}^{\delta} g(u) du \leq \sum_{j=1}^k \frac{\kappa_0^{(j)} A_j (s+1)}{2\delta(2-p)} =: M,$$

where $\kappa_0^{(j)}$ and A_j are the constants from (8) and Lemma 2.2 corresponding to the strip $T_\delta(\gamma^{(j)})$. By the mean-value argument used in the proof of Theorem 1.5, there exists $u_0 \in]-\delta, \delta[\setminus E$ with $g(u_0) \leq M$. In particular, $g_j(u_0) \leq M$ for every j .

Since $u_0 \notin E$, the parallel curve $\gamma_{u_0}^{(j)}$ avoids S for every j . Applying Hölder's inequality (14) with $g_j(u_0) \leq M$ in place of the individual bound, and the optimisation $p = 2 - 1/\log(s+2)$ exactly as in the proof of Theorem 1.5, we obtain

$$\ell_S(\gamma_{u_0}^{(j)}) \leq C'_j \sqrt{(s+1) \log(s+2)}$$

for every $j = 1, \dots, k$, where $C'_j > 0$ depends only on $\alpha_1, \dots, \alpha_k$, B_0 , and ρ .

Set $\hat{\gamma}_j := \gamma_{u_0}^{(j)}$ and $q := q(u_0) = \exp_{b_0}(u_0 \eta_0) \in B_0 \setminus S$. Define the path $\sigma : [0, |u_0|] \rightarrow B_0$ by $\sigma(u) := \exp_{b_0}(\frac{u}{|u_0|} u_0 \eta_0)$ if $u_0 \neq 0$, and let σ be the constant path at b_0 if $u_0 = 0$; it connects b_0 to q with $\ell_\rho(\sigma) = |u_0| \leq \delta$. For each j , the family $\{\gamma_u^{(j)}\}$ for u between 0 and u_0 is a free homotopy in $T_\delta(\gamma^{(j)}) \subset B_0$ from $\gamma_0^{(j)} = \gamma^{(j)}$ (based at b_0) to $\gamma_{u_0}^{(j)} = \hat{\gamma}_j$ (based at q), and this homotopy moves the base point along σ . It follows that $[\sigma^{-1} \circ \hat{\gamma}_j \circ \sigma] = [\gamma^{(j)}] = \alpha_j$ in $\pi_1(B_0, b_0)$ for every j . Finally, $\hat{\gamma}_j = \gamma_{u_0}^{(j)} \subset T_\delta(\gamma^{(j)})$ and $\sigma([0, |u_0|]) \subset T_\delta(\gamma^{(1)})$, so the last assertion holds with $K_0 := \bigcup_{i=1}^k T_\delta(\gamma^{(i)}) \Subset B_0$ and $L_0 := \max_{1 \leq i \leq k} \sup_{|u| \leq \delta} \ell_\rho(\gamma_u^{(i)}) < +\infty$. Setting $C := \max_j C'_j$ and $\delta' := \delta$ completes the proof. $\square$

2.3 Reduction to the simple case

In this subsection we lift the standing simplicity assumption of §2.2 and prove Theorem 1.5 for every nontrivial $\alpha \in \pi_1(B_0, b_0)$.

Since B_0 has the homotopy type of a compact orientable surface of genus g with $t \geq 1$ boundary circles, $\pi_1(B_0, b_0)$ is free of rank $k := 2g + t - 1$ [13, Theorem 3.2.8]. Moreover it admits a basis of smooth simple loops meeting pairwise only at b_0 : model the surface as a closed disk with k attached bands, place b_0 at the center, and let γ_j be the core of the j -th band closed up by the two radii to its endpoints, with the three corners rounded off. The surface deformation-retracts onto $\gamma_1 \cup \dots \cup \gamma_k$, so the classes $\alpha_j := [\gamma_j]$, $j = 1, \dots, k$, form a free basis of $\pi_1(B_0, b_0)$, represented by the smooth simple loops γ_j . For background on curves in surfaces see [3, Chapter 1]. Up to smoothing, such a basis is the *simple base* of [10, §3.1], whose representatives are only piecewise smooth; it is what we call a *simple basis* throughout the paper, in particular in Proposition 2.5.

Proof of Theorem 1.5, general case. Fix a nontrivial loop $\alpha \subset B_0$ and a simple basis $\alpha_1, \dots, \alpha_k$ of $\pi_1(B_0, b_0)$ as above. Write α as a (reduced) word in this basis:

$$\alpha = \alpha_{i_1}^{\epsilon_1} \cdots \alpha_{i_m}^{\epsilon_m} \quad \text{in } \pi_1(B_0, b_0), \quad \epsilon_j \in \{\pm 1\}, \quad i_j \in \{1, \dots, k\}.$$

The word length $m = m(\alpha) \geq 1$ is determined by α and the basis.

Fix $S \subset B_0$ with $\#S = s \geq 1$. By Proposition 2.5 applied to the simple basis $\alpha_1, \dots, \alpha_k$, there exist

- a point $q \in B_0 \setminus S$,
- a path $\sigma: [0, 1] \rightarrow B_0$ from b_0 to q with $\ell_\rho(\sigma) \leq \delta'$,
- loops $\hat{\gamma}_1, \dots, \hat{\gamma}_k \subset B_0 \setminus S$ based at q ,

such that $[\sigma^{-1} \circ \hat{\gamma}_j \circ \sigma] = \alpha_j$ in $\pi_1(B_0, b_0)$ and

$$\ell_S(\hat{\gamma}_j) \leq C_0 \sqrt{(s+1) \log(s+2)} \quad \text{for all } j = 1, \dots, k,$$

where $C_0 > 0$ depends only on $\alpha_1, \dots, \alpha_k$, B_0 , and ρ .

Define the concatenation

$$\hat{\gamma}_\alpha := \hat{\gamma}_{i_m}^{\epsilon_m} \circ \hat{\gamma}_{i_{m-1}}^{\epsilon_{m-1}} \circ \dots \circ \hat{\gamma}_{i_1}^{\epsilon_1},$$

a piecewise smooth loop based at q . We verify that $\hat{\gamma}_\alpha \in \Gamma_S^\alpha$ and bound its Kobayashi length.

Membership in Γ_S^α . Each $\hat{\gamma}_{i_j}$ is contained in $B_0 \setminus S$, hence so is the concatenation $\hat{\gamma}_\alpha$. To see that $\hat{\gamma}_\alpha$ is freely homotopic to α in B_0 , insert $\sigma \circ \sigma^{-1}$ between consecutive factors and use distributivity of conjugation:

$$[\sigma^{-1} \circ \hat{\gamma}_\alpha \circ \sigma] = [\sigma^{-1} \circ \hat{\gamma}_{i_1}^{\epsilon_1} \circ \sigma] \cdots [\sigma^{-1} \circ \hat{\gamma}_{i_m}^{\epsilon_m} \circ \sigma] = \alpha_{i_1}^{\epsilon_1} \cdots \alpha_{i_m}^{\epsilon_m} = \alpha \quad \text{in } \pi_1(B_0, b_0).$$

Thus $\hat{\gamma}_\alpha$, viewed as a loop based at q , is conjugate (via the path σ) to a loop based at b_0 representing α ; equivalently, the free homotopy class of $\hat{\gamma}_\alpha$ in B_0 coincides with the free homotopy class of α .

Kobayashi length bound. Since κ_S is a Finsler pseudo-metric, its homogeneity gives $\kappa_S(b, -v) = \kappa_S(b, v)$, hence $\ell_S(\hat{\gamma}_{i_j}^{-1}) = \ell_S(\hat{\gamma}_{i_j})$. By additivity of ℓ_S over concatenation,

$$\ell_S(\hat{\gamma}_\alpha) = \sum_{j=1}^m \ell_S(\hat{\gamma}_{i_j}^{\epsilon_j}) = \sum_{j=1}^m \ell_S(\hat{\gamma}_{i_j}) \leq m \cdot C_0 \sqrt{(s+1) \log(s+2)}.$$

Conclusion. Setting $C := m \cdot C_0$, which depends only on α (through its word length m in the fixed basis), the simple basis $\alpha_1, \dots, \alpha_k$, B_0 , and ρ , we obtain

$$\ell_S(\Gamma_S^\alpha) \leq \ell_S(\hat{\gamma}_\alpha) \leq C \sqrt{(s+1) \log(s+2)}.$$

As $S \subset B_0$ with $\#S = s \geq 1$ was arbitrary, and since sets with $\#S < s$ satisfy the same bound a fortiori (the right-hand side being increasing in s), taking the supremum over all S with $\#S \leq s$ yields

$$L(\alpha, s) \leq C \sqrt{(s+1) \log(s+2)} \quad \text{for all } s \geq 1.$$

For $s = 0$, $L(\alpha, 0) = \inf_\beta \ell_{\kappa_{B_0}}(\beta)$ is a fixed positive constant, absorbed into C . □

We now record two immediate consequences.

Corollary 2.6. *Fix a loop $\alpha \subset B_0$ representing a nontrivial class in $\pi_1(B_0, b_0)$, then*

$$\lim_{s \rightarrow +\infty} \frac{\log L(\alpha, s)}{\log s} = \frac{1}{2}.$$

Proof. Combining Theorem 1.5 with Phung’s lower bound [10, Theorem C] gives

$$\frac{c\sqrt{s}}{\log(s+2)} \leq L(\alpha, s) \leq C\sqrt{(s+1)\log(s+2)}$$

so the result follows immediately. $\square$

Remark 2.7. Corollary 2.6 gives $L(\alpha, s) = O(s^{1/2+\varepsilon})$ for every $\varepsilon > 0$, but the precise asymptotic behavior of $L(\alpha, s)$ remains open: between Phung’s lower bound $c\sqrt{s}/\log(s+2)$ and our upper bound $C\sqrt{(s+1)\log(s+2)}$ there is still a multiplicative gap of order $(\log(s+2))^{3/2}$. Determining the exact asymptotics of $L(\alpha, s)$ is, to our knowledge, an open problem.

3 Bounding generalized integral points

3.1 Phung’s bound

In this section we revisit the proof of Theorem 1.3. We follow Phung’s strategy [10] closely, and we recall the structure of the argument in enough detail to identify where our improvement enters (§3.2). Our account differs from *loc. cit.* in the following respects; each point is taken up again, with full notation, at the corresponding place of the proof below.

- Phung bounds the Kobayashi–Royden metric of $(\mathcal{A} \setminus \mathcal{D})|_{B_0}$ from below by a fixed hermitian metric by invoking [10, Theorem 7.5]. That statement concerns a relatively compact subset of an arbitrary complex manifold, but its proof is deferred to [4, Theorem 3], which deals with complements of hypersurfaces in complex tori. We give a short self-contained proof of the comparison, in the generality actually needed here, based on Brody’s reparametrization lemma and Hurwitz’s theorem [8, Lemmas 7.2.1 and 7.2.11].
- At one point the argument requires a path of uniformly bounded length joining two points of the same fibre *within that fibre*. In *loc. cit.* this length is bounded by the diameter of the closure of $\mathcal{A}_{B_0}$ in $\mathcal{A}$. This is not sufficient: two points of a fibre may be close to each other in $\mathcal{A}$ even though every path joining them inside the fibre is long. We bound the length instead by the diameter of the fibre itself, measured with respect to the metric induced on the fibre, and we check that these diameters are uniformly bounded.
- Phung converts bounds on the lengths of certain loops in $\mathcal{A}_{B_0}$ into bounds on the displacements, along the fibre direction of the universal cover, of the corresponding deck transformations. This conversion needs justification: the metric pulled back to the universal cover is not a product metric, so the projection onto the fibre direction does not decrease lengths in general. We show that all the lifted paths in question stay in a fixed region, independent of s and of the section, on which this projection is Lipschitz with a uniform constant.
- Several routine reductions are carried out in detail: the case where $\pi_1(B_0)$ is trivial, the reduction to $t \geq 1$, the enlargement of the disks $\overline{\Delta}_i$ absorbing the (countable) set of points of B whose fibre in $\mathcal{D}$ fails to be hyperbolic, and the verification that the two conjugating paths naturally appearing in the construction differ by a loop contained in a fibre, so that the associated cocycles are cohomologous.

The statement and the strategy of Theorem 1.3 are entirely Phung’s; the purpose of the details above is to make the argument self-contained in the form needed for §3.2.

Proof of Theorem 1.3. Fix a hermitian metric h on $\mathcal{A}$. Put $k := \text{rk}(\pi_1(B_0))$, where we have implicitly fixed the base point $b_0 \in B_0$ for $\pi_1(B_0)$.

The rank-zero case. If $k = 0$ then $\pi_1(B_0, b_0)$ is trivial, so $H^1(\pi_1(B_0, b_0), \Gamma) = 0$ for the fibre lattice $\Gamma = H_1(A_{b_0}, \mathbb{Z}) \cong \mathbb{Z}^{2n}$. By the Parshin–Phung injectivity ([9, Proposition 1], [10, Proposition 7.2]; cf. Proposition 3.1 below) the map $A(K)/\text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) \rightarrow H^1(\pi_1(B_0, b_0), \Gamma) = 0$ is the zero homomorphism; hence every two points of $A(K)$ are congruent modulo $\text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) + A(K)_{\text{tors}}$, so $A(K)/\text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C})$ is finite

of order $t_{\mathcal{A}} := \#(A(K)/\mathrm{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}))_{\mathrm{tors}} < \infty$ (Lang–Néron). Consequently $\#(I(s, B_0)/\mathrm{Tr}_{K/\mathbb{C}}(A)(\mathbb{C})) \leq t_{\mathcal{A}}$ for every s , and the bounds of Theorem 1.3 and Theorem 1.4 hold trivially (take $m \geq t_{\mathcal{A}}$). We therefore assume $k \geq 1$ from now on.

Reduction to $t \geq 1$. As in the proof of [10, Theorem A], we may moreover assume $t \geq 1$: if $t = 0$ (so that $\mathrm{Sing}(f) = \emptyset$ and $B_0 = B$ is a compact hyperbolic surface of genus $g \geq 2$), remove one closed disk $\overline{\Delta}_1 \subset B$. This shrinks B_0 , so $I(s, B_0)$ only grows, and it leaves the rank unchanged, $\mathrm{rk} \pi_1(B \setminus \overline{\Delta}_1) = 2g + 1 - 1 = 2g = \mathrm{rk} \pi_1(B)$; hence the bound for the new B_0 implies the bound for the old one. With $t \geq 1$ the surface B_0 has the homotopy type of a compact surface of genus g with $t \geq 1$ boundary circles, so $\pi_1(B_0, b_0)$ is a free group of rank $k = 2g + t - 1$; this freeness will be used in Steps 3 and 6, and it places us within the standing hypotheses of §2.2.

Reduction to $s \geq 1$. Since $I(0, B_0) \subseteq I(1, B_0)$, the bound for $s = 1$ implies the bound for $s = 0$ after enlarging m . We therefore assume $s \geq 1$ from now on.

Step 1: Reduction to the hyperbolic locus. Consider the non-hyperbolic locus

$$V := \{b \in B : \mathcal{D}_b \text{ is not hyperbolic}\}$$

It is well known that V is a closed subset of B in the analytic (Euclidean) topology: the hyperbolic locus of a proper holomorphic map between complex spaces is open ([8, Theorem 7.2.15], applied to $f|_{\mathcal{D}} : \mathcal{D} \rightarrow B$). Moreover, by [10, Lemma B.2], applied via the inclusion $V \subseteq Z(\mathcal{A}, \mathcal{D})$ furnished by [10, Theorem 7.6], and using crucially the assumption (P)(iii) that D contains no translate of a nonzero abelian subvariety, the set V is *at most countable*.

We may therefore enlarge the disks $\overline{\Delta}_1, \dots, \overline{\Delta}_t$ so as to remove $V \cup \mathrm{Sing}(f)$ from the base. First, a collar of each $\partial \overline{\Delta}_i$ is foliated by parallel Jordan curves, and only countably many of these meet the countable set $V \cup \mathrm{Sing}(f)$; we may thus enlarge each $\overline{\Delta}_i$ inside its collar to a closed disk $\overline{\Delta}_i^+$ whose boundary avoids $V \cup \mathrm{Sing}(f)$, the $\overline{\Delta}_i^+$ remaining pairwise disjoint. In particular $\mathrm{Sing}(f) \subset \bigsqcup_i \mathrm{int} \overline{\Delta}_i^+$. The set $W := (V \cup \mathrm{Sing}(f)) \setminus \bigsqcup_i \mathrm{int} \overline{\Delta}_i^+$ is closed and disjoint from the compact set $\bigsqcup_i \overline{\Delta}_i^+$, hence at positive ρ -distance d_0 from it. Applying [10, Lemma 8.1] to W with disks of radius at most ε , for some $0 < \varepsilon < \min\{d_0/2, \frac{1}{3} \min_{i \neq j} d_\rho(\overline{\Delta}_i^+, \overline{\Delta}_j^+)\}$, yields finitely many pairwise disjoint closed cover disks whose union contains W in its interior and each of which is disjoint from every $\overline{\Delta}_j^+$. Annex each cover disk V' to some $\overline{\Delta}_i^+$ by setting $\overline{\Delta}_i' := \overline{\Delta}_i^+ \cup \beta \cup V'$, where β is a thin tube running from $\overline{\Delta}_i^+$ to V' , meeting each only in a short boundary arc at its respective end, and disjoint from all other disks and tubes. Each $\overline{\Delta}_i'$ is simply connected with piecewise- C^1 boundary, hence again a closed disk (in general not a metric ball), and the $\overline{\Delta}_i'$ are pairwise disjoint, with distinct points of $\mathrm{Sing}(f)$ still lying in distinct disks. By construction $V \cup \mathrm{Sing}(f) \subseteq \bigsqcup_{i=1}^t \mathrm{int} \overline{\Delta}_i'$. Writing $B'_0 := B \setminus \bigsqcup_{i=1}^t \overline{\Delta}_i' \subseteq B_0$, the surface B'_0 is a deformation retract of B_0 (one pushes the attached collars, tubes and cover disks out through $\partial \overline{\Delta}_i'$); in particular $\pi_1(B'_0) = \pi_1(B_0)$, so $\mathrm{rk} \pi_1(B'_0) = \mathrm{rk} \pi_1(B_0) = k$ is unchanged. Since $B'_0 \subseteq B_0$ we have $I(s, B_0) \subseteq I(s, B'_0)$ for every s , so it suffices to prove the bound for B'_0 . Replacing B_0 by B'_0 , we may thus assume from now on that $V \cup \mathrm{Sing}(f) \subseteq \bigsqcup_{i=1}^t \mathrm{int} \overline{\Delta}_i$; equivalently, the compact set $\overline{B_0}$ is disjoint from $V \cup \mathrm{Sing}(f)$. In particular $\mathcal{D}_b$ is Kobayashi hyperbolic for every $b \in \overline{B_0}$; hence $\mathcal{D}_b$ contains no translate of a positive-dimensional subtorus of $\mathcal{A}_b$, since such a translate would be a non-hyperbolic complex subspace of the hyperbolic space $\mathcal{D}_b$. By a result of Green ([8, Theorem 7.3.8(ii)], applied to the underlying reduced divisor of $\mathcal{D}_b$; see also [10, Theorem 7.6]) it follows that $\mathcal{A}_b \setminus \mathcal{D}_b$ is hyperbolically embedded into $\mathcal{A}_b$, in particular Kobayashi hyperbolic, for any $b \in \overline{B_0}$. Finally, fix a compact connected smooth subsurface $N \Subset B_0$ with $b_0 \in N$ (it will be enlarged below finitely many times, by fixed compact sets independent of s and P). Let δ_N be the *intrinsic* diameter of N , i.e. the diameter of N for the length metric induced by ρ on N ; it is finite, and it is realized by paths in N , since N with this length metric is a compact geodesic space. Hence for any two points $x, y \in N$ we can find a path $c_{x,y} \subset N$ joining them and such that:

$$\ell_\rho(c_{x,y}) \leq \delta_N < +\infty.$$

We denote $\mathcal{C}_{b_0} := \{c_{b_0,p}\}_{p \in N}$.

Step 2: Bounding loop lengths in the base. Let $\sigma_P \in I(s, B_0)$ and set $S := f(\sigma_P(B_0) \cap \mathcal{D}) \cap B_0$, so that $\#S \leq s$ and $\sigma_P(B_0 \setminus S) \subseteq (\mathcal{A} \setminus \mathcal{D})|_{B_0 \setminus S}$; we may assume $S \neq \emptyset$ (otherwise replace S by an arbitrary singleton of B_0 : this only increases the Kobayashi lengths below, and $\#S \leq s$ still holds since $s \geq 1$). Let $\alpha_1, \dots, \alpha_k$ be a simple basis of $\pi_1(B_0, b_0)$. By [10, Theorem 5.1], applied with $\mathcal{C}_{b_0}$ as the fixed collection of paths of [10, Definition 3.2] (possible since the base points b produced there lie in $B_\varepsilon \subseteq N$, cf. below),

there exist $b \in B_0$ and piecewise smooth loops $\gamma_j \in \Gamma_S^{\alpha_j}$ such that $\alpha_j = c_{b_0,b}^{-1} \gamma_j c_{b_0,b}$ in $\pi_1(B_0, b_0)$ where $c_{b_0,b} \in \mathcal{C}_{b_0}$ and the following bound holds:

$$\ell_S(\gamma_j) \leq L(s+1) \quad \forall j = 1, \dots, k \quad (19)$$

for $L \in \mathbb{R}_{>0}$ independent of s, S, b . Two further facts, immediate from the proof of [10, Theorem 5.1], will be needed in Step 5: the loops γ_j lie in a fixed compact set $\mathcal{K}_1 \Subset B_0$ and satisfy $\ell_\rho(\gamma_j) \leq L_1$, with $\mathcal{K}_1$ and L_1 independent of s, S, P . Indeed, by [10, Lemmas 3.7 and 4.1] the γ_j are obtained by modifying, within ρ -distance $2a$, finitely many fixed loops of bounded ρ -length, while keeping ρ -distance $\geq 2a$ from ∂B_0 ; the length bound follows from [10, (5.2)–(5.3)]. Moreover, the base points b lie in the fixed compact set $B_\varepsilon \Subset B_0$ of *loc. cit.* We enlarge N once and for all so that $\mathcal{K}_1 \cup B_\varepsilon \subseteq N$.

There exists $c \in \mathbb{R}_{>0}$ such that, at the infinitesimal level,

$$\kappa_{(\mathcal{A} \setminus \mathcal{D})|_{B_0}}(x, v) \geq c|v|_h \quad \forall (x, v). \quad (20)$$

This is [10, Theorem 7.5], a criterion of Brody–Green type going back to [4, Theorem 3]; as the proof is only indicated in *loc. cit.*, we give the short argument. If no such c exists then, by the definition of the Kobayashi–Royden metric (§2.1), there are holomorphic disks $g_\nu: \Delta(0, 1) \rightarrow (\mathcal{A} \setminus \mathcal{D})|_{B_0}$ with $|g'_\nu(0)|_h \rightarrow \infty$. Since $(\mathcal{A} \setminus \mathcal{D})|_{B_0}$ is relatively compact in the complex manifold $\mathcal{A}|_{B \setminus \text{Sing}(f)}$ (its closure lies over $\overline{B_0}$, which is disjoint from $\text{Sing}(f)$ by the reduction above), Brody’s reparametrization lemma [8, Lemma 7.2.1] produces a non-constant entire curve $F: \mathbb{C} \rightarrow \mathcal{A}$ which is a locally uniform limit of holomorphic maps with values in $(\mathcal{A} \setminus \mathcal{D})|_{B_0}$. In particular $f \circ F(\mathbb{C}) \subseteq \overline{B_0}$ omits the nonempty open set $\text{int } \overline{\Delta_1}$, hence omits three points of B ; so $f \circ F$ maps into the complement of three points of B , which is hyperbolic by uniformization, and $f \circ F$ is therefore constant. Hence $F(\mathbb{C}) \subseteq \mathcal{A}_b$ for some $b \in \overline{B_0}$, and $b \notin V \cup \text{Sing}(f)$ by the reduction above. By Hurwitz’s theorem in the form [8, Lemma 7.2.11], applied on an exhaustion of $\mathbb{C}$ by disks inside the complex manifold $\mathcal{A}|_{B \setminus \text{Sing}(f)}$, where $\mathcal{D}$ restricts to a reduced divisor, either $F(\mathbb{C}) \subseteq \mathcal{D}_b$ or $F(\mathbb{C}) \subseteq \mathcal{A}_b \setminus \mathcal{D}_b$; both are Kobayashi hyperbolic since $b \notin V$ (Step 1), so F is constant — a contradiction.

Now notice that

$$\sigma_P(\gamma_j) \subset \sigma_P(B_0 \setminus S) \subseteq (\mathcal{A} \setminus \mathcal{D})|_{B_0 \setminus S}$$

So by Equations (19) to (20) and the decreasing property of the Kobayashi–Royden metric we deduce

$$\ell_h(\sigma_P(\gamma_j)) \leq c^{-1}L(s+1). \quad (21)$$

Step 3: The Parshin cocycle. Since $f: \mathcal{A}_{B_0} \rightarrow B_0$ is a proper smooth submersion, Ehresmann’s theorem gives a fiber bundle in the differentiable category. Let σ_O be the zero section and fix $w_0 := \sigma_O(b_0) \in \mathcal{A}_{b_0}$. Since B_0 is a $K(\pi, 1)$ -space (i.e. $\pi_i(B_0) = 0$ for $i \geq 2$, which holds because its universal cover is the disc $\mathbb{D}$), the long exact sequence of homotopy groups for the fibration $\mathcal{A}_{b_0} \rightarrow \mathcal{A}_{B_0} \rightarrow B_0$ yields the short exact sequence

$$1 \longrightarrow \pi_1(\mathcal{A}_{b_0}, w_0) \longrightarrow \pi_1(\mathcal{A}_{B_0}, w_0) \xrightarrow{f_*} \pi_1(B_0, b_0) \longrightarrow 1. \quad (22)$$

As $\mathcal{A}_{b_0}$ is a complex torus of dimension n , we have $\pi_1(\mathcal{A}_{b_0}, w_0) = H_1(\mathcal{A}_{b_0}, \mathbb{Z}) =: \Gamma \cong \mathbb{Z}^{2n}$. The zero section σ_O splits (22) via $i_O(\alpha) := [\sigma_O(\gamma)]$ where $\alpha = [\gamma] \in G := \pi_1(B_0, b_0)$. Each rational point $P \in A(K)$ with section $\sigma_P: B_0 \rightarrow \mathcal{A}_{B_0}$ gives a second splitting

$$i_P(\alpha) := [\eta_{w_0, \sigma_P(b_0)}^{-1} \circ \sigma_P(\gamma) \circ \eta_{w_0, \sigma_P(b_0)}],$$

where $\eta_{w_0, \sigma_P(b_0)}$ is a path in the fiber $\mathcal{A}_{b_0}$ from w_0 to $\sigma_P(b_0)$. Since both i_P and i_O are sections of f_* , for every $\alpha \in G$:

$$f_*(i_P(\alpha) \cdot i_O(\alpha)^{-1}) = \alpha \cdot \alpha^{-1} = 1,$$

so $i_P(\alpha) \cdot i_O(\alpha)^{-1} \in \ker(f_*) = \Gamma$. We define the *Parshin cocycle*

$$c_P: G \rightarrow \Gamma, \quad c_P(\alpha) := i_P(\alpha) \cdot i_O(\alpha)^{-1}.$$

One verifies that c_P is a 1-cocycle of G with coefficients in the G -module Γ , where the G -action on Γ is by conjugation

$$\begin{aligned} \varphi: G &\rightarrow \text{Aut}(\Gamma) \\ \alpha &\mapsto \varphi_\alpha: \gamma \mapsto i_O(\alpha) \gamma i_O(\alpha)^{-1} \end{aligned}$$

Note that the action φ doesn't depend on the choice of the splitting i_O , in fact for another splitting i_P , we have $i_P(\alpha) = c_P(\alpha) \cdot i_O(\alpha)$ with $c_P(\alpha) \in \Gamma$, and

$$i_P(\alpha) \gamma i_P(\alpha)^{-1} = c_P(\alpha) \cdot \varphi_\alpha(\gamma) \cdot c_P(\alpha)^{-1} = \varphi_\alpha(\gamma),$$

where the last equality uses the commutativity of Γ . The cohomology class $[c_P] \in H^1(G, \Gamma)$ is independent of the choice of path $\eta_{w_0, \sigma_P(b_0)}$ (a different choice changes c_P by a coboundary). The following result proved in [9, Proposition 1] and [10, Proposition 7.2] will be crucial:

Proposition 3.1 (Parshin-Phung). *The homomorphism*

$$\begin{aligned} \Psi: A(K) &\rightarrow H^1(G, \Gamma) \\ P &\mapsto [c_P] \end{aligned}$$

factors through $A(K)/\text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C})$ and has the following injectivity property: if $\Psi(P) = \Psi(Q)$, then $P - Q \in \text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) + A(K)_{\text{tors}}$. In particular, each element $[c] \in H^1(G, \Gamma)$ in the image of Ψ arises from at most $t_{\mathcal{A}} := \#(A(K)/\text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}))_{\text{tors}} < \infty$ rational points modulo the trace (finite by the Lang-Néron theorem).

Remark 3.2. It would be interesting to understand the relationship between the Parshin cocycle and the “analytic cocycle” induced by the logarithm of the section σ_P , as defined in [2, Equation (13)].

To extract a quantitative bound from the cocycle c_P , we work in the semidirect product coordinates provided by i_O . The splitting i_O exhibits

$$\pi_1(\mathcal{A}_{B_0}, w_0) \cong \Gamma \rtimes_{\varphi} G. \quad (23)$$

In these coordinates, $i_O(\alpha_j) = (0, \alpha_j)$ and $i_P(\alpha_j) = (\beta_j, \alpha_j)$ for some $\beta_j \in \Gamma$, so the Parshin cocycle reads $c_P(\alpha_j) = \beta_j$. Since G is free, the k -tuple $(\beta_1, \dots, \beta_k) \in \Gamma^k$ determines c_P completely. The displacement bound in Step 4 will give a norm bound on each β_j .

Step 4: Bounding loop lengths in $\mathcal{A}$. Let us now go back to the special loops (based in b) $\gamma_j \in \Gamma_S^{\alpha_j}$ for $j = 1, \dots, k$ considered in Step 2. Consider the following loop based at w_0 :

$$\sigma_P(\gamma_j)^{\#} := (v_b \circ \sigma_O(c_{b_0, b}))^{-1} \circ \sigma_P(\gamma_j) \circ (v_b \circ \sigma_O(c_{b_0, b}))$$

where $v_b \subset \mathcal{A}_b$ is a h -geodesic from $\sigma_O(b)$ to $\sigma_P(b)$. Since $\sigma_P(\gamma_j) = \sigma_P(c_{b_0, b}) \circ \sigma_P(\alpha_j) \circ \sigma_P(c_{b_0, b})^{-1}$, $\sigma_P(\gamma_j)^{\#}$ is freely homotopic to $\sigma_P(\alpha_j)$ and each of its components satisfies the following uniform bounds:

- the loop $\sigma_P(\gamma_j)$ satisfies:

$$\ell_h(\sigma_P(\gamma_j)) \leq c^{-1}L(s+1)$$

by Equation (21);

- the fiber path $v_b \subset \mathcal{A}_b$ satisfies:

$$\ell_h(v_b) \leq \delta_0 := \sup_{b \in \overline{B_0}} \text{diam}_h(\mathcal{A}_b) < +\infty;$$

(here $\text{diam}_h(\mathcal{A}_b)$ is the *intrinsic* h -diameter of the fibre, which bounds the length of a fibre geodesic; the supremum is finite because $\overline{B_0} \cap \text{Sing}(f) = \emptyset$ by Step 1, so $(\mathcal{A}_b)_{b \in \overline{B_0}}$ is a compact smooth family and $b \mapsto \text{diam}_h(\mathcal{A}_b)$ is continuous)

- since B is compact $\|d\sigma_O\|_{\infty} := \sup_{b \in B} \|d_b \sigma_O\| < +\infty$, so

$$\ell_h(\sigma_O(c_{b_0, b})) \leq \|d\sigma_O\|_{\infty} \ell(c_{b_0, b}) \leq \|d\sigma_O\|_{\infty} \delta_N =: \delta'_0$$

In other words for any element of the simple basis α_j we can construct a loop $[\sigma_P(\gamma_j)^{\#}] \in \pi_1(\mathcal{A}_{B_0}, w_0)$ that is freely homotopic to $\sigma_P(\alpha_j)$ and satisfies

$$\ell_h(\sigma_P(\gamma_j)^{\#}) \leq H(s) := c^{-1}L(s+1) + 2(\delta_0 + \delta'_0)$$

where the constants L, δ_0, δ'_0 don't depend on S and P .

The loop $\sigma_P(\gamma_j)^\#$ is based at w_0 , so it defines an element $[\sigma_P(\gamma_j)^\#] \in \pi_1(\mathcal{A}_{B_0}, w_0)$. This element need not equal $i_P(\alpha_j) = (\beta_j, \alpha_j)$, because $\sigma_P(\gamma_j)^\#$ and the loop defining $i_P(\alpha_j)$ use different conjugating paths from w_0 to the fibre above b : the former uses $v_b \circ \sigma_O(c_{b_0, b})$, while the latter uses $\sigma_P(c_{b_0, b}) \circ \eta_{w_0, \sigma_P(b_0)}$. Both are paths from w_0 to $\sigma_P(b)$, so their concatenation

$$\omega := (v_b \circ \sigma_O(c_{b_0, b}))^{-1} \circ \sigma_P(c_{b_0, b}) \circ \eta_{w_0, \sigma_P(b_0)}$$

is a loop at w_0 with $[\omega] \in \Gamma$: indeed $f_*[\omega] = 1$, since $f \circ v_b$ and $f \circ \eta_{w_0, \sigma_P(b_0)}$ are constant while $f \circ \sigma_O(c_{b_0, b}) = f \circ \sigma_P(c_{b_0, b}) = c_{b_0, b}$. A direct verification gives

$$[\sigma_P(\gamma_j)^\#] = [\omega]^{-1} \cdot i_P(\alpha_j) \cdot [\omega] \quad \text{in } \pi_1(\mathcal{A}_{B_0}, w_0).$$

In other words, $[\sigma_P(\gamma_j)^\#]$ and $i_P(\alpha_j)$ are *conjugate* in $\pi_1(\mathcal{A}_{B_0}, w_0)$. The map $i'_P: G \rightarrow \pi_1(\mathcal{A}_{B_0}, w_0)$ defined by $i'_P(\alpha_j) := [\sigma_P(\gamma_j)^\#]$ is therefore a section of (22) (conjugation by $[\omega] \in \Gamma = \ker f_*$ preserves the section property), Γ -conjugate to i_P , so the associated cocycles are cohomologous: $[c'] = [c_P] \in H^1(G, \Gamma)$. By Proposition 3.1, we may freely replace i_P by i'_P for counting purposes. We do so and, to lighten notation, simply write β_j for the components of the new cocycle.

Step 5: Displacement bound via the universal cover. We now explain how the h -length bound on $\sigma_P(\gamma_j)^\#$ constrains the lattice element $\beta_j \in \Gamma$. Let $\pi: \tilde{\mathcal{A}}_0 \rightarrow \mathcal{A}_{B_0}$ be the universal cover. Since $\mathcal{A}_{B_0} \rightarrow B_0$ is a fibre bundle (by Ehresmann's theorem) with fibre $\mathcal{A}_{b_0}$ (a complex torus of dimension n , so $\mathcal{A}_{b_0} \cong \mathbb{R}^{2n}$) and base B_0 (a hyperbolic Riemann surface, so $\tilde{B}_0 \cong \mathbb{D}$, the unit disk), pulling back to $\mathbb{D}$ trivialises the bundle and gives $\tilde{\mathcal{A}}_0 \cong \mathbb{R}^{2n} \times \mathbb{D}$. We use the trivialisation induced by the relative exponential (cf. [10, (7.3)]): the lattice local system $(R^1 f_* \mathbb{Z})^\vee$ is trivial over the contractible $\mathbb{D}$, and with this choice the subgroup $\Gamma \subset \pi_1(\mathcal{A}_{B_0}, w_0)$ acts on $\mathbb{R}^{2n} \times \mathbb{D}$ by $(\tilde{x}, \tilde{y}) \mapsto (\tilde{x} + \gamma, \tilde{y})$.

Fix the point $\tilde{w} = (\tilde{x}_0, \tilde{y}_0) \in \pi^{-1}(w_0)$ with $\tilde{x}_0 = 0$: this choice is possible because $w_0 = \sigma_O(b_0)$ lies on the zero section, whose lift in the chosen trivialisation is $\{0\} \times \mathbb{D}$, and $\tilde{y}_0$ is any point of $\mathbb{D}$ lying over b_0 for the universal covering $v: \mathbb{D} \rightarrow B_0$. Pull back the hermitian metric h to a Riemannian metric $\tilde{h}$ on $\tilde{\mathcal{A}}_0$. The group of deck transformations of π is $\pi_1(\mathcal{A}_{B_0}, w_0) \cong \Gamma \rtimes_\varphi G$ (with the composition convention for π_1 already implicit in our loop concatenations). The action of an element $(\beta, \alpha) = \beta \cdot i_O(\alpha) \in \Gamma \rtimes_\varphi G$ on $\mathbb{R}^{2n} \times \mathbb{D}$ is given by (homotopy lifting property, cf. the proof of [10, Theorem A]):

$$(\beta, \alpha) \cdot (\tilde{x}, \tilde{y}) = (\varphi_\alpha(\tilde{x}) + \beta, \alpha \cdot \tilde{y}), \quad (24)$$

where $\alpha \cdot \tilde{y}$ is the deck transformation of $v: \mathbb{D} \rightarrow B_0$ by $\alpha \in G = \pi_1(B_0, b_0)$ (acting on the base), and φ_α acts on the fibre factor $\mathbb{R}^{2n}$ as the linear extension of the monodromy: the fibre component “twists” as one transports along the base loop α , and the semidirect product structure of $\Gamma \rtimes_\varphi G$ encodes precisely this twisting. In particular the fibre component of the action is a translation only when $\varphi_\alpha = \text{id}$; at the chosen point $\tilde{w} = (\tilde{x}_0, \tilde{y}_0)$, however, thanks to the normalisation $\tilde{x}_0 = 0$, (24) reads

$$(\beta, \alpha) \cdot (\tilde{x}_0, \tilde{y}_0) = (\beta \cdot \tilde{x}_0, \alpha \cdot \tilde{y}_0), \quad (25)$$

where $\beta \cdot \tilde{x}_0$ is the deck transformation of $u: \mathbb{R}^{2n} \rightarrow \mathcal{A}_{b_0}$ by the element $\beta \in \Gamma$ (acting on the fibre).

The loop $\sigma_P(\gamma_j)^\#$ is based at w_0 and has homotopy class (β_j, α_j) . Its lift to $\tilde{\mathcal{A}}_0$ from $\tilde{w}$ is a path from $\tilde{w}$ to $(\beta_j, \alpha_j) \cdot \tilde{w}$. Since π is a local isometry for the metrics $\tilde{h}$ and h , the $\tilde{h}$ -length of the lift equals $\ell_h(\sigma_P(\gamma_j)^\#) \leq H(s)$, so:

$$d_{\tilde{h}}(\tilde{w}, (\beta_j, \alpha_j) \cdot \tilde{w}) \leq H(s) := c^{-1}L(s+1) + 2(\delta_0 + \delta'_0) = O(s), \quad (26)$$

where L, δ_0, δ'_0 are independent of S and P . Let d_j be the Γ -invariant geodesic metric on the fibre factor $\mathbb{R}^{2n}$ obtained by restricting $\tilde{h}$ to the slice $\mathbb{R}^{2n} \times \{\alpha_j \cdot \tilde{y}_0\}$; its quotient by the translation action of $\Gamma \cong \mathbb{Z}^{2n}$ is the compact torus $\mathcal{A}_{b_0}$. Let $\text{pr}_1: \mathbb{R}^{2n} \times \mathbb{D} \rightarrow \mathbb{R}^{2n}$ be the projection; by (25), $\text{pr}_1(\tilde{w}) = \tilde{x}_0$ and $\text{pr}_1((\beta_j, \alpha_j) \cdot \tilde{w}) = \beta_j \cdot \tilde{x}_0$.

The $\mathbb{D}$ -component of the lift $\tilde{c}$ of $\sigma_P(\gamma_j)^\#$ from $\tilde{w}$ is the lift, from $\tilde{y}_0$, of the base loop $f(\sigma_P(\gamma_j)^\#) = c_{b_0, b}^{-1} \circ \gamma_j \circ c_{b_0, b}$, which by Step 2 has image in N and ρ -length at most $L_2 := L_1 + 2\delta_N$, uniformly in s, P . Hence this component is a path of $\tilde{\rho}$ -length $\leq L_2$ starting at $\tilde{y}_0$ and contained in $v^{-1}(N)$, so it remains in the closed ball

$$\Omega := \overline{B}_{v^{-1}(N)}(\tilde{y}_0, L_2)$$

of radius L_2 around $\tilde{y}_0$ in $v^{-1}(N)$ for the induced length metric. The space $(v^{-1}(N), \tilde{\rho})$ is a covering of the compact length space N with the lifted length metric, hence complete and locally compact, hence a

proper metric space by the Hopf–Rinow theorem for length spaces; therefore Ω is compact. Note that Ω is independent of s and P , and $\tilde{c} \subset \mathbb{R}^{2n} \times \Omega$.

The ratio $|\text{pr}_{1*}\xi|_{d_j}/|\xi|_{\tilde{h}}$, for $0 \neq \xi \in T(\mathbb{R}^{2n} \times \Omega)$, is invariant under Γ : every $\gamma \in \Gamma$ is a $\tilde{h}$ -isometry and satisfies $\text{pr}_1 \circ \gamma = \tau_\gamma \circ \text{pr}_1$, with τ_γ the d_j -isometric translation by γ . (The analogous equivariance fails for a general deck element (β, α) , whose fibre component has linear part φ_α by (24); this is why we restrict to the Γ -invariant region $\mathbb{R}^{2n} \times \Omega$.) The ratio therefore descends to a continuous function on the unit sphere bundle over the compact set $\mathcal{A}_{b_0} \times \Omega$, hence it is bounded by some $C_{h,j} \geq 1$; we set $C_h := \max_{1 \leq j \leq k} C_{h,j}$, which depends only on $\mathcal{A}$, h , Ω and the basis, uniformly in s, P . Therefore pr_1 is C_h -Lipschitz from $(\mathbb{R}^{2n} \times \Omega, \tilde{h})$ to $(\mathbb{R}^{2n}, d_j)$, so $\text{pr}_1 \circ \tilde{c}$ has d_j -length $\leq C_h \ell_{\tilde{h}}(\tilde{c})$, and

$$d_j(\beta_j \cdot \tilde{x}_0, \tilde{x}_0) \leq C_h \ell_{\tilde{h}}(\tilde{c}) = C_h \ell_h(\sigma_P(\gamma_j)^\#) \leq C_h H(s). \quad (27)$$

Remark 3.3. The intermediate bound (26) controls the displacement in the *full* universal cover $\tilde{\mathcal{A}}_0 \cong \mathbb{R}^{2n} \times \mathbb{D}$, involving both fibre and base directions (and the metric $\tilde{h}$ is not a product metric due to the monodromy). The projection step (27) extracts a purely fibre-theoretic bound, which is what we need for the lattice count in the next step.

Step 6: Counting. For each $j = 1, \dots, k$, we count the elements $\beta \in \Gamma$ compatible with (27):

$$N_j(s) := \#\left\{\beta \in \Gamma : d_j(\beta \cdot \tilde{x}_0, \tilde{x}_0) \leq C_h H(s)\right\}. \quad (28)$$

Via the universal covering $u: \mathbb{R}^{2n} \rightarrow \mathcal{A}_{b_0}$, the torus A_{b_0} becomes a compact geodesic Riemannian manifold with the metric d_j , and $\Gamma \cong \mathbb{Z}^{2n}$ acts on $(\mathbb{R}^{2n}, d_j)$ by deck transformations. The set in (28) is in natural bijection with the set of Γ -translates of $\tilde{x}_0$ lying in the d_j -ball of radius $C_h H(s)$ around $\tilde{x}_0$.

By the fundamental lemma of the geometry of groups (cf. [10, Proposition A.8 and Lemma A.9]), since $\Gamma \cong \mathbb{Z}^{2n}$ is a lattice of rank $2n$ acting cocompactly on $(\mathbb{R}^{2n}, d_j)$, there exists $m_j > 0$ (depending on $\mathcal{A}$, h , α_j , but not on s) such that

$$N_j(s) \leq m_j (C_h H(s) + 1)^{2n}. \quad (29)$$

The exponent $2n$ reflects the rank of $\Gamma \cong \mathbb{Z}^{2n}$: in a ball of radius R in the universal cover of the $2n$ -dimensional real torus $\mathcal{A}_{b_0}$, the number of Γ -translates of $\tilde{x}_0$ grows as R^{2n} .

Since G is free on $\alpha_1, \dots, \alpha_k$, the cohomology class $[c_P] \in H^1(G, \Gamma)$ is completely determined by the k -tuple $(\beta_1, \dots, \beta_k) \in \Gamma^k$. Bounding each component independently via (29) with $R = C_h H(s)$:

$$\#\{\text{possible cohomology classes } [c_P]\} \leq \prod_{j=1}^k m_j (C_h H(s) + 1)^{2n} = m_0 (C_h H(s) + 1)^{2nk}, \quad (30)$$

where $m_0 := \prod_{j=1}^k m_j$.

By Proposition 3.1, each class $[c_P] \in H^1(G, \Gamma)$ arises from at most $t_A := \#(A(K)/\text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}))_{\text{tors}} < \infty$ rational points modulo the trace (finite by the Lang–Néron theorem). Therefore:

$$\# \left(I(s, B_0) / \text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) \right) \leq t_A \cdot m_0 (C_h H(s) + 1)^{2nk} \leq m (s + 1)^{2nk},$$

where $m := t_A \cdot m_0 \cdot (C_h(c^{-1}L + 2(\delta_0 + \delta'_0)) + 1)^{2nk}$ depends only on B_0 , $\mathcal{A}$, D , and h . $\square$

3.2 Improved bound

We now prove a sharper version of Theorem 1.3. The general strategy is unchanged: we follow the same six-step scheme (reduction to the hyperbolic locus, loop construction, Parshin cocycle, lifting to $\mathcal{A}$, displacement in the universal cover, lattice counting). The only modification occurs in Steps 2 and 4, where we replace Phung’s linear bound $\ell_S(\gamma_j) = O(s)$ with the sublinear bound $O(\sqrt{(s+1)\log(s+2)})$ provided by Proposition 2.5. This propagates through the counting argument and halves the exponent.

Proof of Theorem 1.4. The proof follows the same six-step scheme as the proof of Theorem 1.3; in particular we may again assume $k \geq 1$ and $s \geq 1$, by the same reductions as there. The main modification is in Step 2: we replace the appeal to [10, Theorem 5.1], which provides loops of Kobayashi length $O(s)$, with Proposition 2.5, applied to the set $S := f(\sigma_P(B_0) \cap \mathcal{D}) \cap B_0$ (as in Step 2, we may assume $S \neq \emptyset$; note $\#S \leq s$, and since the bound below is increasing in the cardinality we may state it with s in place of $\#S$), which provides a point $b \in B_0 \setminus S$, a path σ from b_0 to b in B_0 with $\ell_\rho(\sigma) \leq \delta'$, and loops $\gamma_1, \dots, \gamma_k \subset B_0 \setminus S$ based at b with $[\sigma^{-1} \circ \gamma_j \circ \sigma] = \alpha_j$ in $\pi_1(B_0, b_0)$ and

$$\ell_S(\gamma_j) \leq C\sqrt{(s+1)\log(s+2)} \quad \text{for all } j = 1, \dots, k.$$

Since the loops share a common base point b and the conjugation uses the single path σ , Steps 3–5 apply verbatim with the conjugating path $v_b \circ \sigma_O(\sigma)$ in $\mathcal{A}_{B_0}$ (in place of $v_b \circ \sigma_O(c_{b_0, b})$ in Theorem 1.3), whose h -length is bounded by $\delta_0 + \|d\sigma_O\|_\infty \delta'$, independently of s . The compactness input of Step 5 is again available: by the last assertion of Proposition 2.5, the base loop $\sigma^{-1} \circ \gamma_j \circ \sigma$ has image in the fixed compact set $\mathcal{K}_0 \Subset B_0$ and ρ -length at most $L_0 + 2\delta'$, uniformly in s, P ; enlarging N so that $\mathcal{K}_0 \subseteq N$, the compact set Ω of Step 5 (with $L_2 := L_0 + 2\delta'$) is independent of s, P . The displacement bound becomes

$$d_j(\beta_j \cdot \tilde{x}_0, \tilde{x}_0) \leq C_h H(s), \quad H(s) := c^{-1} C \sqrt{(s+1)\log(s+2)} + 2(\delta_0 + \|d\sigma_O\|_\infty \delta'),$$

with $C_h \geq 1$ the constant of Step 5. The lattice counting in Step 6 then gives

$$\# \left(I(s, B_0) / \text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) \right) \leq t_{\mathcal{A}} m_0 (C_h H(s) + 1)^{2nk}.$$

It remains to extract the exponent. Since $C_h \geq 1$, for s large enough

$$C_h H(s) + 1 \leq C_h c^{-1} (C + 1) \sqrt{(s+1)\log(s+2)},$$

hence $(C_h H(s) + 1)^{2n} \leq (C_h c^{-1} (C + 1))^{2n} ((s+1)\log(s+2))^n$. Fix $\varepsilon' > 0$. Since $\log(s+2) \leq (s+1)^{\varepsilon'/n}$ for s sufficiently large, there exists $s_0 = s_0(\varepsilon') \geq 0$ such that for all $s \geq s_0$:

$$((s+1)\log(s+2))^n \leq (s+1)^{n+\varepsilon'}.$$

Over k basis elements, $(C_h H(s) + 1)^{2nk} \leq \text{const} \cdot (s+1)^{nk+k\varepsilon'}$ for $s \geq s_0$. Setting $\varepsilon := k\varepsilon'$ and absorbing all constants into m (for the finitely many values $s < s_0$ use $I(s, B_0) \subseteq I(s_0, B_0)$ and enlarge m), we obtain

$$\# \left(I(s, B_0) / \text{Tr}_{K/\mathbb{C}}(A)(\mathbb{C}) \right) \leq m (s+1)^{nk+\varepsilon} \quad \text{for all } s \geq 0,$$

where $m = m(B_0, \mathcal{A}, \mathcal{D}, h, \varepsilon) > 0$ is independent of s . □

References

- [1] D. Borthwick. *Spectral theory of infinite-area hyperbolic surfaces*, volume 318 of *Progress in Mathematics*. Birkhäuser/Springer, second edition, 2016.
- [2] P. Dolce and F. Tropeano. Relative monodromy of ramified sections on abelian schemes. *International Mathematics Research Notices*, 2026(4):rnag025, 02 2026.
- [3] B. Farb and D. Margalit. *A Primer on Mapping Class Groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [4] M. L. Green. Holomorphic maps to complex tori. *Amer. J. Math.*, 100(3):615–620, 1978.
- [5] M. Hindry and J. H. Silverman. The canonical height and integral points on elliptic curves. *Invent. Math.*, 93(2):419–450, 1988.
- [6] J.M. Lee. *Introduction to Riemannian manifolds*, volume 176 of *Graduate Texts in Mathematics*. Springer, Cham, 2018.
- [7] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2 edition, 2001.

- [8] J. Noguchi and J. Winkelmann. *Nevanlinna Theory in Several Complex Variables and Diophantine Approximation*, volume 350 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2014.
- [9] A. N. Parshin. Finiteness theorems and hyperbolic manifolds. In *The Grothendieck Festschrift, Vol. III*, volume 88 of *Progr. Math.*, pages 163–178. Birkhäuser Boston, Boston, MA, 1990.
- [10] X. K. Phung. Generalized integral points on abelian varieties and the geometric Lang-Vojta conjecture. *J. Algebraic Geom.*, 34(3):407–446, 2025.
- [11] H. L. Royden. Remarks on the Kobayashi metric. In *Several complex variables, II (Proc. Internat. Conf., Univ. Maryland, College Park, Md., 1970)*, Lecture Notes in Math., Vol. 185, pages 125–137. Springer, Berlin-New York, 1971.
- [12] T. Shioda. On elliptic modular surfaces. *J. Math. Soc. Japan*, 24:20–59, 1972.
- [13] H. Zieschang, E. Vogt, and H.-D. Coldewey. *Surfaces and planar discontinuous groups*, volume 835 of *Lecture Notes in Mathematics*. Springer, Berlin, 1980. Translated from the German by J. Stillwell.

P. Dolce, INSTITUTE FOR THEORETICAL SCIENCES, WESTLAKE UNIVERSITY, CHINA
E-mail address: dolce@westlake.edu.cn